

1227292

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office

*September 18, 2004*

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/495,637  
FILING DATE: *August 15, 2003*  
RELATED PCT APPLICATION NUMBER: *PCT/US04/26012*

Certified by



Jon W Dudas

Acting Under Secretary of Commerce  
for Intellectual Property  
and Acting Director of the U.S.  
Patent and Trademark Office

## PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a **PROVISIONAL APPLICATION FOR PATENT** under 37 CFR 1.53 (c).

		DOCKET NUMBER	21481PV
INVENTOR(S)			
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)	
Paul J. Christopher D. Robert M. George D.	Coleman Cox Garbaccio Hartman	Wallingford, Pennsylvania Harleysville, Pennsylvania Lansdale, Pennsylvania Lansdale, Pennsylvania	
<input type="checkbox"/> Additional inventors are being named on the separately numbered sheets attached hereto			
TITLE OF THE INVENTION (500 characters max)			
MITOTIC KINESIN INHIBITORS			
CORRESPONDENCE ADDRESS			
Direct all Correspondence to: Merck & Co., Inc. Patent Department - RY60-30 P.O. Box 2000 Rahway			
		<input checked="" type="checkbox"/> Customer Number	000210
STATE	New Jersey	ZIP CODE	07065
		COUNTRY	U.S.A.
ENCLOSED APPLICATION PARTS (check all that apply)			
<input checked="" type="checkbox"/> Specification	Number of Pages	171	<input type="checkbox"/> CD(s), Number
<input type="checkbox"/> Drawing(s)	Number of Sheets		<input type="checkbox"/> Other (specify)
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76			
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)			
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees		FILING FEE AMOUNT (\$)	\$160.00
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 13-2755			

19587 U.S. PTO  
60/495637  
08/15/03


The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are: \_\_\_\_\_

Respectfully submitted,

SIGNATURE



Date 08/15/2003

TYPED or PRINTED NAME David A. Muthard

REGISTRATION NO. 35,297  
(if appropriate)

TELEPHONE (732) 594-3903

NOTE: Mail to Mail Stop Provisional Application

EXPRESS MAIL CERTIFICATE	
DATE OF DEPOSIT	August 15, 2003
EXPRESS MAIL NO.	EV321986216US
I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL "POST OFFICE TO ADDRESSEE" ON THE ABOVE DATE IN AN ENVELOPE ADDRESSED TO COMMISSIONER FOR PATENTS, P.O. BOX 1450, ALEXANDRIA, VA 22313-1450.	
MAILED BY	DATE August 15, 2003

TITLE OF THE INVENTION  
MITOTIC KINESIN INHIBITORS

5 BACKGROUND OF THE INVENTION

This invention relates to 2,2-disubstituted 2,5-dihydropyrrole derivatives that are inhibitors of mitotic kinesins, in particular the mitotic kinesin KSP, and are useful in the treatment of cellular proliferative diseases, for example cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders and inflammation.

10 Among the therapeutic agents used to treat cancer are the taxanes and vinca alkaloids. Taxanes and vinca alkaloids act on microtubules, which are present in a variety of cellular structures. Microtubules are the primary structural element of the mitotic spindle. The mitotic spindle is responsible for distribution of replicate copies of the genome to each of the two daughter cells that result from cell division. It is presumed that disruption of the mitotic spindle by these drugs results in inhibition of cancer cell division, and induction of cancer cell death.  
15 However, microtubules form other types of cellular structures, including tracks for intracellular transport in nerve processes. Because these agents do not specifically target mitotic spindles, they have side effects that limit their usefulness.

20 Improvements in the specificity of agents used to treat cancer is of considerable interest because of the therapeutic benefits which would be realized if the side effects associated with the administration of these agents could be reduced. Traditionally, dramatic improvements in the treatment of cancer are associated with identification of therapeutic agents acting through novel mechanisms. Examples of this include not only the taxanes, but also the camptothecin class of topoisomerase I inhibitors. From both of these perspectives, mitotic kinesins are  
25 attractive targets for new anti-cancer agents.

Mitotic kinesins are enzymes essential for assembly and function of the mitotic spindle, but are not generally part of other microtubule structures, such as in nerve processes. Mitotic kinesins play essential roles during all phases of mitosis. These enzymes are "molecular motors" that transform energy released by hydrolysis of ATP into mechanical force which drives  
30 the directional movement of cellular cargoes along microtubules. The catalytic domain sufficient for this task is a compact structure of approximately 340 amino acids. During mitosis, kinesins organize microtubules into the bipolar structure that is the mitotic spindle. Kinesins mediate movement of chromosomes along spindle microtubules, as well as structural changes in the mitotic spindle associated with specific phases of mitosis. Experimental perturbation of mitotic

kinesin function causes malformation or dysfunction of the mitotic spindle, frequently resulting in cell cycle arrest and cell death.

Among the mitotic kinesins which have been identified is KSP. KSP belongs to an evolutionarily conserved kinesin subfamily of plus end-directed microtubule motors that assemble into bipolar homotetramers consisting of antiparallel homodimers. During mitosis KSP associates with microtubules of the mitotic spindle. Microinjection of antibodies directed against KSP into human cells prevents spindle pole separation during prometaphase, giving rise to monopolar spindles and causing mitotic arrest and induction of programmed cell death. KSP and related kinesins in other, non-human, organisms, bundle antiparallel microtubules and slide them relative to one another, thus forcing the two spindle poles apart. KSP may also mediate in anaphase B spindle elongation and focussing of microtubules at the spindle pole.

Human KSP (also termed HsEg5) has been described [Blangy, et al., Cell, 83:1159-69 (1995); Whitehead, et al., Arthritis Rheum., 39:1635-42 (1996); Galgio et al., J. Cell Biol., 135:339-414 (1996); Blangy, et al., J Biol. Chem., 272:19418-24 (1997); Blangy, et al., Cell Motil Cytoskeleton, 40:174-82 (1998); Whitehead and Rattner, J. Cell Sci., 111:2551-61 (1998); Kaiser, et al., JBC 274:18925-31 (1999); GenBank accession numbers: X85137, NM004523 and U37426] , and a fragment of the KSP gene (TRIP5) has been described [Lee, et al., Mol Endocrinol., 9:243-54 (1995); GenBank accession number L40372]. Xenopus KSP homologs (Eg5), as well as Drosophila K-LP61 F/KRP 130 have been reported.

Certain quinazolinones have recently been described as being inhibitors of KSP (PCT Publ. WO 01/30768, May 3, 2001).


Mitotic kinesins are attractive targets for the discovery and development of novel mitotic chemotherapeutics. Accordingly, it is an object of the present invention to provide compounds, methods and compositions useful in the inhibition of KSP, a mitotic kinesin.

## SUMMARY OF THE INVENTION

The present invention relates to dihydropyrrole derivatives, that are useful for treating cellular proliferative diseases, for treating disorders associated with KSP kinesin activity, and for inhibiting KSP kinesin. The compounds of the invention may be illustrated by the


Formula I:





# DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are useful in the inhibition of mitotic kinesins and are illustrated by a compound of Formula I:



5

or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

- 10    a is    0 or 1;  
       b is    0 or 1;

m is 0, 1, or 2;  
 n is 0, 1, 2 or 3;  
 r is 0 or 1;  
 s is 0 or 1;  
 5 t is 0, 1 or 2;  
 u is 0 or 1;

R<sup>1</sup> and R<sup>2</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

10

R<sup>3</sup> is selected from:

- 1) hydrogen;
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl;
- 3) C<sub>1</sub>-C<sub>10</sub> alkyl-O-R<sup>d</sup>,
- 15 4) C<sub>2</sub>-C<sub>10</sub> alkenyl-O-R<sup>d</sup>,
- 5) C<sub>2</sub>-C<sub>10</sub> alkynyl-O-R<sup>d</sup>,
- 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-O-R<sup>d</sup>,
- 7) C<sub>1</sub>-C<sub>10</sub> alkyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
- 8) C<sub>2</sub>-C<sub>10</sub> alkenyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
- 20 9) C<sub>2</sub>-C<sub>10</sub> alkynyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
- 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
- 11) C<sub>1</sub>-C<sub>10</sub> alkyl-S(O)<sub>m</sub>-R<sup>d</sup>,
- 12) C<sub>2</sub>-C<sub>10</sub> alkenyl-S(O)<sub>m</sub>-R<sup>d</sup>,
- 13) C<sub>2</sub>-C<sub>10</sub> alkynyl-S(O)<sub>m</sub>-R<sup>d</sup>,
- 25 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-S(O)<sub>m</sub>-R<sup>d</sup>,

said alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more substituents selected from R<sup>6</sup>;

30 R<sup>4</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) CO<sub>2</sub>H,
- 4) halo,
- 35 5) CN,

- 5
- 6) OH,
  - 7)  $O_bC_1-C_6$  perfluoroalkyl,
  - 8)  $O_a(C=O)_bNR^8R^9$ ,
  - 9)  $S(O)_mR^a$ ,
  - 10)  $S(O)_2NR^8R^9$ , and
  - 11)  $-OPO(OH)_2$ ;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

10

R<sup>5</sup> is selected from:

- 15
- 1) hydrogen;
  - 2)  $(C=O)_aO_bC_1-C_{10}$  alkyl,
  - 3)  $(C=O)_aO_b$ aryl,
  - 4)  $CO_2H$ ,
  - 5) halo,
  - 6) CN,
  - 7) OH,
  - 8)  $O_bC_1-C_6$  perfluoroalkyl,
  - 20 9)  $O_a(C=O)_bNR^8R^9$ ,
  - 10)  $S(O)_mR^a$ ,
  - 11)  $S(O)_2NR^8R^9$ , and
  - 12)  $-OPO(OH)_2$ ;

25 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>6</sup> is independently selected from:

- 30
- 1)  $(C=O)_aO_bC_1-C_{10}$  alkyl,
  - 2)  $(C=O)_aO_b$ aryl,
  - 3)  $C_2-C_{10}$  alkenyl,
  - 4)  $C_2-C_{10}$  alkynyl,
  - 5)  $(C=O)_aO_b$  heterocyclyl,
  - 6)  $CO_2H$ ,
  - 35 7) halo,

- 8) CN,  
 9) OH,  
 10)  $O_bC_1-C_6$  perfluoroalkyl,  
 11)  $O_a(C=O)_bNR^8R^9$ ,  
 5 12)  $S(O)_mR^a$ ,  
 13)  $S(O)_2NR^8R^9$ ,  
 14) oxo,  
 15) CHO,  
 16)  $(N=O)R^8R^9$ ,  
 10 17)  $(C=O)_aO_bC_3-C_8$  cycloalkyl, and  
 18)  $-OPO(OH)_2$ ;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from  $R^7$ ;

15

$R^7$  is selected from:

- 1)  $(C=O)_rO_s(C_1-C_{10})$ alkyl,  
 2)  $O_r(C_1-C_3)$ perfluoroalkyl,  
 3) oxo,  
 20 4) OH,  
 5) halo,  
 6) CN,  
 7)  $(C_2-C_{10})$ alkenyl,  
 8)  $(C_2-C_{10})$ alkynyl,  
 25 9)  $(C=O)_rO_s(C_3-C_6)$ cycloalkyl,  
 10)  $(C=O)_rO_s(C_0-C_6)$ alkylene-aryl,  
 11)  $(C=O)_rO_s(C_0-C_6)$ alkylene-heterocyclyl,  
 12)  $(C=O)_rO_s(C_0-C_6)$ alkylene- $N(R^b)_2$ ,  
 13)  $C(O)R^a$ ,  
 30 14)  $(C_0-C_6)$ alkylene- $CO_2R^a$ ,  
 15)  $C(O)H$ ,  
 16)  $(C_0-C_6)$ alkylene- $CO_2H$ , and  
 17)  $(C=O)_rN(R^b)_2$ ,  
 18)  $S(O)_mR^a$ ,  
 35 19)  $S(O)_2N(R^b)_2$ ; and

20)  $-\text{OPO}(\text{OH})_2$ ;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $\text{R}^b$ , OH,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , halogen,  $\text{CO}_2\text{H}$ , CN,

5  $\text{O}(\text{C}=\text{O})\text{C}_1\text{-C}_6$  alkyl, oxo,  $\text{NO}_2$  and  $\text{N}(\text{R}^b)_2$ ;

$\text{R}^8$  and  $\text{R}^9$  are independently selected from:

- 1) H,
- 2)  $(\text{C}=\text{O})\text{O}_b\text{C}_1\text{-C}_{10}$  alkyl,
- 10 3)  $(\text{C}=\text{O})\text{O}_b\text{C}_3\text{-C}_8$  cycloalkyl,
- 4)  $(\text{C}=\text{O})\text{O}_b\text{aryl}$ ,
- 5)  $(\text{C}=\text{O})\text{O}_b\text{heterocyclyl}$ ,
- 6)  $\text{C}_1\text{-C}_{10}$  alkyl,
- 7) aryl,
- 15 8)  $\text{C}_2\text{-C}_{10}$  alkenyl,
- 9)  $\text{C}_2\text{-C}_{10}$  alkynyl,
- 10) heterocyclyl,
- 11)  $\text{C}_3\text{-C}_8$  cycloalkyl,
- 12)  $\text{SO}_2\text{R}^a$ , and
- 20 13)  $(\text{C}=\text{O})\text{NR}^b_2$ ,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from  $\text{R}^7$ , or

$\text{R}^8$  and  $\text{R}^9$  can be taken together with the nitrogen to which they are attached to form a  
 25 monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ;

30  $\text{R}^{10}$  and  $\text{R}^{11}$  are independently selected from: F and  $-\text{CH}_2\text{F}$ ;

$\text{R}^{12}$  and  $\text{R}^{13}$  are independently selected from: H and  $-\text{CH}_2\text{F}$ ;

$\text{R}^{\text{ox}}$  is absent or is oxo;

35



$R^a$  is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from  $R^7$ ;

$R^b$  is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup> or S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one, two or three substituents selected from  $R^7$ ;

$R^c$  and  $R^{c'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, NH<sub>2</sub>, OH, OR<sup>a</sup>, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup>, S(O)<sub>2</sub>R<sup>a</sup> and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N(R<sup>b</sup>)<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from  $R^7$ ; or


$R^c$  and  $R^{c'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^7$ ;

$R^d$  is selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, -(C<sub>2</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N(R<sup>b</sup>)<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from  $R^7$ ;

$R^e$  and  $R^{e'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from  $R^7$ ; or

$R^e$  and  $R^{e'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^7$ .

The compounds of this invention are useful in the inhibition of mitotic kinesins and are illustrated by a compound of Formula II:



or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

- 5       a is    0 or 1;  
        b is    0 or 1;  
        m is    0, 1, or 2;  
        n is    0, 1, 2 or 3;  
 10       r is    0 or 1;  
        s is    0 or 1;  
        t is    0 or 1;
- R<sup>1</sup> and R<sup>2</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-  
 15 C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>3</sup> is selected from:

- 20       1)    hydrogen;  
        2)    C<sub>1</sub>-C<sub>10</sub> alkyl;  
        3)    C<sub>1</sub>-C<sub>10</sub> alkyl-O-R<sup>d</sup>,  
        4)    C<sub>2</sub>-C<sub>10</sub> alkenyl-O-R<sup>d</sup>,  
        5)    C<sub>2</sub>-C<sub>10</sub> alkynyl-O-R<sup>d</sup>,  
        6)    (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-O-R<sup>d</sup>,

- 7) C<sub>1</sub>-C<sub>10</sub> alkyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
  - 8) C<sub>2</sub>-C<sub>10</sub> alkenyl-(C=O)<sub>b</sub>NR<sup>c</sup>R<sup>c'</sup>,
  - 9) C<sub>2</sub>-C<sub>10</sub> alkynyl-(C=O)<sub>b</sub>NR<sup>c</sup>R<sup>c'</sup>,
  - 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-(C=O)<sub>b</sub>NR<sup>c</sup>R<sup>c'</sup>,
  - 5 11) C<sub>1</sub>-C<sub>10</sub> alkyl-S(O)<sub>m</sub>-R<sup>d</sup>,
  - 12) C<sub>2</sub>-C<sub>10</sub> alkenyl- S(O)<sub>m</sub>-R<sup>d</sup>,
  - 13) C<sub>2</sub>-C<sub>10</sub> alkynyl- S(O)<sub>m</sub>-R<sup>d</sup>,
  - 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl- S(O)<sub>m</sub>-R<sup>d</sup>,
- 10 said alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more substituents selected from R<sup>6</sup>;

R<sup>4</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 15 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) CO<sub>2</sub>H,
- 4) halo,
- 5) CN,
- 6) OH,
- 20 7) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 8) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>8</sup>R<sup>9</sup>,
- 9) S(O)<sub>m</sub>R<sup>a</sup>,
- 10) S(O)<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

- 25 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>5</sup> is selected from:

- 1) hydrogen;
- 2) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 30 3) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 4) CO<sub>2</sub>H,
- 5) halo,
- 6) CN,
- 7) OH,

- 8)  $O_bC_1-C_6$  perfluoroalkyl,
- 9)  $O_a(C=O)_bNR^8R^9$ ,
- 10)  $S(O)_mR^a$ ,
- 11)  $S(O)_2NR^8R^9$ , and
- 5 12)  $-OPO(OH)_2$ ;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from  $R^7$ ;

10  $R^6$  is independently selected from:

- 1)  $(C=O)_aO_bC_1-C_{10}$  alkyl,
- 2)  $(C=O)_aO_b$  aryl,
- 3)  $C_2-C_{10}$  alkenyl,
- 4)  $C_2-C_{10}$  alkynyl,
- 15 5)  $(C=O)_aO_b$  heterocyclyl,
- 6)  $CO_2H$ ,
- 7) halo,
- 8)  $CN$ ,
- 9)  $OH$ ,
- 20 10)  $O_bC_1-C_6$  perfluoroalkyl,
- 11)  $O_a(C=O)_bNR^8R^9$ ,
- 12)  $S(O)_mR^a$ ,
- 13)  $S(O)_2NR^8R^9$ ,
- 14) oxo,
- 25 15)  $CHO$ ,
- 16)  $(N=O)R^8R^9$ ,
- 17)  $(C=O)_aO_bC_3-C_8$  cycloalkyl, and
- 18)  $-OPO(OH)_2$ ;

30 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from  $R^7$ ;

$R^7$  is selected from:

- 1)  $(C=O)_rO_s(C_1-C_{10})$  alkyl,
- 35 2)  $O_r(C_1-C_3)$  perfluoroalkyl,

- 3) oxo,  
 4) OH,  
 5) halo,  
 6) CN,  
 5 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,  
 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,  
 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,  
 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,  
 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,  
 10 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>,  
 13) C(O)R<sup>a</sup>,  
 14) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>,  
 15) C(O)H,  
 16) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H,  
 15 17) C(O)N(R<sup>b</sup>)<sub>2</sub>,  
 18) S(O)<sub>m</sub>R<sup>a</sup>,  
 19) S(O)<sub>2</sub>N(R<sup>b</sup>)<sub>2</sub>; and  
 20) -OPO(OH)<sub>2</sub>;

20 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R<sup>b</sup>, OH, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, CO<sub>2</sub>H, CN, O(C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, oxo, NO<sub>2</sub> and N(R<sup>b</sup>)<sub>2</sub>;

R<sup>8</sup> and R<sup>9</sup> are independently selected from:

- 25 1) H,  
 2) (C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,  
 3) (C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 4) (C=O)O<sub>b</sub>aryl,  
 5) (C=O)O<sub>b</sub>heterocyclyl,  
 30 6) C<sub>1</sub>-C<sub>10</sub> alkyl,  
 7) aryl,  
 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,  
 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,  
 10) heterocyclyl,  
 35 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,



12)  $\text{SO}_2\text{R}^a$ , and

13)  $(\text{C}=\text{O})\text{NR}^b_2$ ,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from  $\text{R}^7$ , or

5

$\text{R}^8$  and  $\text{R}^9$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ;

10

$\text{R}^{10}$  is selected from: F and  $-\text{CH}_2\text{F}$ ;

$\text{R}^{13}$  is selected from: H and  $-\text{CH}_2\text{F}$ , provided that if t is 1,  $\text{R}^{13}$  is H;

15

$\text{R}^{\text{ox}}$  is absent or is oxo;

$\text{R}^a$  is independently selected from:  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$ , aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ;

20

$\text{R}^b$  is independently selected from: H,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ , aryl, heterocyclyl,  $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$ ,  $(\text{C}=\text{O})\text{OC}_1\text{-C}_6\text{ alkyl}$ ,  $(\text{C}=\text{O})\text{C}_1\text{-C}_6\text{ alkyl}$ ,  $(\text{C}=\text{O})\text{aryl}$ ,  $(\text{C}=\text{O})\text{heterocyclyl}$ ,  $(\text{C}=\text{O})\text{NR}^e\text{R}^e$  or  $\text{S}(\text{O})_2\text{R}^a$ , optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ;

25

$\text{R}^c$  and  $\text{R}^{c'}$  are independently selected from: H,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ , aryl,  $\text{NH}_2$ , OH,  $\text{OR}^a$ ,  $-(\text{C}_1\text{-C}_6)\text{alkyl-OH}$ ,  $-(\text{C}_1\text{-C}_6)\text{alkyl-O}-(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}=\text{O})\text{OC}_1\text{-C}_6\text{ alkyl}$ ,  $(\text{C}=\text{O})\text{C}_1\text{-C}_6\text{ alkyl}$ ,  $(\text{C}=\text{O})\text{aryl}$ ,  $(\text{C}=\text{O})\text{heterocyclyl}$ ,  $(\text{C}=\text{O})\text{NR}^e\text{R}^e$ ,  $\text{S}(\text{O})_2\text{R}^a$  and  $-(\text{C}_1\text{-C}_6)\text{alkyl-N}(\text{R}^b)_2$ , wherein the alkyl is optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ; or

30

$\text{R}^c$  and  $\text{R}^{c'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ;


35

$R^d$  is selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, -(C<sub>2</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N(R<sup>b</sup>)<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

$R^e$  and  $R^{e'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or

$R^e$  and  $R^{e'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>.

In an embodiment of the invention the compounds are illustrated by a compound of Formula III:



or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

a is 0 or 1;  
 b is 0 or 1;  
 m is 0, 1, or 2;  
 n is 0, 1 or 2;

r is 0 or 1;

s is 0 or 1;

R<sup>1</sup> and R<sup>2</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,

5 optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>4</sup> is independently selected from:

- 1) halo,
- 2) OH,
- 10 3) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

R<sup>5</sup> is selected from:

- 1) hydrogen,
- 2) halo,
- 15 3) OH,
- 4) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

R<sup>7</sup> is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 20 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 3) oxo,
- 4) OH,
- 5) halo,
- 6) CN,
- 25 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,
- 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,
- 30 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>,
- 13) C(O)R<sup>a</sup>,
- 14) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>,
- 15) C(O)H,
- 16) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H, and
- 35 17) C(O)N(R<sup>b</sup>)<sub>2</sub>,

18)  $S(O)_m R^a$ , and

19)  $S(O)_2 N(R^b)_2$ ;

5 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $R^b$ , OH,  $(C_1-C_6)$ alkoxy, halogen,  $CO_2H$ , CN,  $O(C=O)C_1-C_6$  alkyl, oxo,  $NO_2$  and  $N(R^b)_2$ ;

$R^8$  and  $R^9$  are independently selected from:

- 1) H,
- 10 2)  $(C=O)O_b C_1-C_{10}$  alkyl,
- 3)  $(C=O)O_b C_3-C_8$  cycloalkyl,
- 4)  $(C=O)O_b$ aryl,
- 5)  $(C=O)O_b$ heterocyclyl,
- 6)  $C_1-C_{10}$  alkyl,
- 15 7) aryl,
- 8)  $C_2-C_{10}$  alkenyl,
- 9)  $C_2-C_{10}$  alkynyl,
- 10) heterocyclyl,
- 11)  $C_3-C_8$  cycloalkyl,
- 20 12)  $SO_2 R^a$ , and
- 13)  $(C=O)NR^b_2$ ,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from  $R^7$ , or

25  $R^8$  and  $R^9$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^7$ ;

30  $R^a$  is independently selected from:  $(C_1-C_6)$ alkyl,  $(C_3-C_6)$ cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from  $R^7$ ;

$R^b$  is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup> or S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

5    R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, NH<sub>2</sub>, OH, OR<sup>a</sup>, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup>, S(O)<sub>2</sub>R<sup>a</sup> and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N(R<sup>b</sup>)<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or


10   R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

15   R<sup>e</sup> and R<sup>e'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or

20   R<sup>e</sup> and R<sup>e'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>.

25                    Another embodiment of the present invention is illustrated by a compound of Formula IV:





or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

5

a is 0 or 1;

b is 0 or 1;

m is 0, 1, or 2;

r is 0 or 1;

10 s is 0 or 1;

**R<sup>1</sup>** and **R<sup>2</sup>** are independently selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl, optionally substituted with one, two or three substituents selected from **R<sup>7</sup>**;

15 **R<sup>4</sup>** is independently selected from:

- 1) halo,
- 2) OH,
- 3) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

20 **R<sup>7</sup>** is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 3) oxo,
- 4) OH,
- 25 5) halo,

- 6) CN,  
 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,  
 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,  
 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,  
 5 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,  
 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,  
 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>,  
 13) C(O)R<sup>a</sup>,  
 14) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>,  
 10 15) C(O)H,  
 16) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H, and  
 17) C(O)N(R<sup>b</sup>)<sub>2</sub>,  
 18) S(O)<sub>m</sub>R<sup>a</sup>, and  
 19) S(O)<sub>2</sub>N(R<sup>b</sup>)<sub>2</sub>;

15

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R<sup>b</sup>, OH, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, CO<sub>2</sub>H, CN, O(C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, oxo, NO<sub>2</sub> and N(R<sup>b</sup>)<sub>2</sub>;

20 R<sup>8</sup> and R<sup>9</sup> are independently selected from:

- 1) H,  
 2) (C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,  
 3) (C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 4) (C=O)O<sub>b</sub>aryl,  
 25 5) (C=O)O<sub>b</sub>heterocyclyl,  
 6) C<sub>1</sub>-C<sub>10</sub> alkyl,  
 7) aryl,  
 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,  
 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,  
 30 10) heterocyclyl,  
 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 12) SO<sub>2</sub>R<sup>a</sup>, and  
 13) (C=O)NR<sup>b</sup><sub>2</sub>,

35 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>, or

R<sup>8</sup> and R<sup>9</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said  
 5 monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>a</sup> is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

10 R<sup>b</sup> is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NReRe' or S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

15 R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, NH<sub>2</sub>, OH, OR<sup>a</sup>, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NReRe', S(O)<sub>2</sub>R<sup>a</sup> and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N(R<sup>b</sup>)<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or


20 R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

25 Re and Re' are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or

30 Re and Re' can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>.

A further embodiment of the present invention is illustrated by a compound of

35 Formula V:




or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

5

$R^1$  and  $R^2$  are independently selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

A further embodiment of the present invention is illustrated by a compound of Formula VI:



10

or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

15  $R^1$  and  $R^2$  are independently selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

Specific examples of the compounds of the instant invention include:

- 5 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 10 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 15 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 20 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(2*R*,4*R*)-2-(fluoromethyl)-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 25 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(2*S*,4*S*)-2-(fluoromethyl)-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 30 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methyl-1-oxidopiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoropiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 35




(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-isopropylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide


5 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*S*)-3-fluoropiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

or a pharmaceutically acceptable salt or stereoisomer thereof.

10 Particular examples of the compounds of the instant invention are:




15 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide




(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

5



(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(2*R*,4*R*)-2-(fluoromethyl)-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide



(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

5 or a pharmaceutically acceptable salt or stereoisomer thereof.

The compounds of the present invention may have asymmetric centers, chiral axes, and chiral planes (as described in: E.L. Eliel and S.H. Wilen, *Stereochemistry of Carbon Compounds*, John Wiley & Sons, New York, 1994, pages 1119-1190), and occur as racemates,  
 10 racemic mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers, all such stereoisomers being included in the present invention. In addition, the compounds disclosed herein may exist as tautomers and both tautomeric forms are intended to be encompassed by the scope of the invention, even though only one tautomeric structure is depicted.

15 When any variable (e.g. R<sup>4</sup>, R<sup>7</sup>, R<sup>10</sup>, etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every other occurrence. Also, combinations of substituents and variables are permissible only if such combinations result in stable compounds. Lines drawn into the ring systems from substituents represent that the indicated bond may be attached to any of the substitutable ring atoms. If the ring system is  
 20 polycyclic, it is intended that the bond be attached to any of the suitable carbon atoms on the proximal ring only.

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well  
 25 as those methods set forth below, from readily available starting materials. If a substituent is

itself substituted with more than one group, it is understood that these multiple groups may be on the same carbon or on different carbons, so long as a stable structure results. The phrase “optionally substituted with one or more substituents” should be taken to be equivalent to the phrase “optionally substituted with at least one substituent” and in such cases the preferred embodiment will have from zero to three substituents.

As used herein, “alkyl” is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, C<sub>1</sub>-C<sub>10</sub>, as in “C<sub>1</sub>-C<sub>10</sub> alkyl” is defined to include groups having 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbons in a linear or branched arrangement. For example, “C<sub>1</sub>-C<sub>10</sub> alkyl” specifically includes methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *t*-butyl, *i*-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, and so on. The term “cycloalkyl” means a monocyclic saturated aliphatic hydrocarbon group having the specified number of carbon atoms. For example, “cycloalkyl” includes cyclopropyl, methyl-cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, cyclohexyl, and so on. In an embodiment of the invention the term “cycloalkyl” includes the groups described immediately above and further includes monocyclic unsaturated aliphatic hydrocarbon groups. For example, “cycloalkyl” as defined in this embodiment includes cyclopropyl, methyl-cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, cyclohexyl, cyclopentenyl, cyclobutenyl and so on.

The term “alkylene” means a hydrocarbon diradical group having the specified number of carbon atoms. For example, “alkylene” includes -CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>- and the like.

When used in the phrases “C<sub>1</sub>-C<sub>6</sub> aralkyl” and “C<sub>1</sub>-C<sub>6</sub> heteroaralkyl” the term “C<sub>1</sub>-C<sub>6</sub>” refers to the alkyl portion of the moiety and does not describe the number of atoms in the aryl and heteroaryl portion of the moiety.

“Alkoxy” represents either a cyclic or non-cyclic alkyl group of indicated number of carbon atoms attached through an oxygen bridge. “Alkoxy” therefore encompasses the definitions of alkyl and cycloalkyl above.

If no number of carbon atoms is specified, the term “alkenyl” refers to a non-aromatic hydrocarbon radical, straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to four non-aromatic carbon-carbon double bonds may be present. Thus, “C<sub>2</sub>-C<sub>6</sub> alkenyl” means an alkenyl radical having from 2 to 6 carbon atoms. Alkenyl groups include ethenyl, propenyl, butenyl, 2-methylbutenyl and cyclohexenyl. The straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated.

The term "alkynyl" refers to a hydrocarbon radical straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond. Up to three carbon-carbon triple bonds may be present. Thus, "C<sub>2</sub>-C<sub>6</sub> alkynyl" means an alkynyl radical having from 2 to 6 carbon atoms. Alkynyl groups include ethynyl, propynyl, butynyl, 3-methylbutynyl and so on. The straight, branched or cyclic portion of the alkynyl group may contain triple bonds and may be substituted if a substituted alkynyl group is indicated.


In certain instances, substituents may be defined with a range of carbons that includes zero, such as (C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl. If aryl is taken to be phenyl, this definition would include phenyl itself as well as -CH<sub>2</sub>Ph, -CH<sub>2</sub>CH<sub>2</sub>Ph, CH(CH<sub>3</sub>)CH<sub>2</sub>CH(CH<sub>3</sub>)Ph, and so on.

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydronaphthyl, indanyl and biphenyl. In cases where the aryl substituent is bicyclic and one ring is non-aromatic, it is understood that attachment is via the aromatic ring.


The term heteroaryl, as used herein, represents a stable monocyclic or bicyclic ring of up to 7 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Heteroaryl groups within the scope of this definition include but are not limited to: acridinyl, carbazolyl, cinnolinyl, quinoxalinyl, pyrazolyl, indolyl, benzotriazolyl, furanyl, thienyl, benzothienyl, benzofuranyl, quinolinyl, isoquinolinyl, oxazolyl, isoxazolyl, indolyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, tetrahydroquinoline. As with the definition of heterocycle below, "heteroaryl" is also understood to include the N-oxide derivative of any nitrogen-containing heteroaryl. In cases where the heteroaryl substituent is bicyclic and one ring is non-aromatic or contains no heteroatoms, it is understood that attachment is via the aromatic ring or via the heteroatom containing ring, respectively.

The term "heterocycle" or "heterocyclyl" as used herein is intended to mean a 5- to 10-membered aromatic or nonaromatic heterocycle containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S, and includes bicyclic groups. "Heterocyclyl" therefore includes the above mentioned heteroaryls, as well as dihydro and tetrathydro analogs thereof. Further examples of "heterocyclyl" include, but are not limited to the following: benzoimidazolyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, carbolinyl, cinnolinyl, furanyl, imidazolyl, indolinyl, indolyl, indolazinyl, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinazolinyl,

- quinolyl, quinoxaliny, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydroisoquinoliny, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, azetidiny, 1,4-dioxanyl, hexahydroazepiny, piperaziny, piperidiny, pyridin-2-ony, pyrrolidiny, morpholiny, thiomorpholiny, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyraziny, dihydropyrazolyl, dihydropyridiny, dihydropyrimidiny, dihydropyrroly, dihydroquinoliny, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl, dihydroazetidiny, methylenedioxybenzoyl, tetrahydrofuranyl, and tetrahydrothienyl, and N-oxides thereof.
- 10 Attachment of a heterocyclyl substituent can occur via a carbon atom or via a heteroatom.
- Preferably, heterocycle is selected from 2-azepinone, benzimidazolyl, 2-diazapinone, imidazolyl, 2-imidazolidinone, indolyl, isoquinoliny, morpholiny, piperidyl, piperaziny, pyridyl, pyrrolidiny, 2-piperidinone, 2-pyrimidinone, 2-pyrrolidinone, quinoliny, tetrahydrofuryl, tetrahydroisoquinoliny, and thienyl.
- 15 As appreciated by those of skill in the art, "halo" or "halogen" as used herein is intended to include chloro, fluoro, bromo and iodo.
- The alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl substituents may be substituted or unsubstituted, unless specifically defined otherwise. For example, a (C<sub>1</sub>-C<sub>6</sub>)alkyl may be substituted with one, two or three substituents selected from
- 20 OH, oxo, halogen, alkoxy, dialkylamino, or heterocyclyl, such as morpholiny, piperidiny, and so on. In this case, if one substituent is oxo and the other is OH, the following are included in the definition:
- C(=O)CH<sub>2</sub>CH(OH)CH<sub>3</sub>, -(C=O)OH, -CH<sub>2</sub>(OH)CH<sub>2</sub>CH(O), and so on.
- In certain instances, R<sup>c</sup> and R<sup>c'</sup> and R<sup>f</sup> and R<sup>f'</sup> are defined such that they can be
- 25 taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said heterocycle optionally substituted with one or more substituents selected from R<sup>7</sup>. Examples of the heterocycles that can thus be formed include, but are not limited to the following, keeping in mind that the
- 30 heterocycle is optionally substituted with one or more (and in an embodiment, one, two or three) substituents chosen from R<sup>7</sup>:



In certain instances,  $R^d$  and  $R^{d'}$  are defined such that they can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members in the ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from  $N^{Re}$ , O and S, said heterocycle optionally substituted with one or more substituents selected from  $R^7$ . Examples of the heterocycles that can thus be formed include, but are not limited to the following, keeping in mind that the heterocycle is optionally substituted with one or more (and in an embodiment, one or two) substituents chosen from  $R^7$ :



In an embodiment,  $R^1$  is selected from H and  $C_1$ - $C_6$  alkyl.

In an embodiment,  $R^2$  is selected from H and  $C_1$ - $C_6$  alkyl.

In an embodiment,  $R^{12}$  is H.

5 In an embodiment,  $R^3$  is selected from  $-C_1$ - $C_{10}$  alkyl-O- $R^8$  and  $-C_1$ - $C_{10}$  alkyl- $NR^fR^f$ , optionally substituted with one to two substituents selected from  $R^{10}$ .

In an embodiment,  $R^4$  is independently selected from halogen and OH. In a further embodiment, n is 2 and  $R^4$  is independently selected from halogen.

In an embodiment,  $R^5$  is independently selected from H, halogen and OH.

10 In an embodiment, u is 0.

In an embodiment, t is 1,  $R^{10}$  is fluoro and  $R^{13}$  is H.

In another embodiment, t is 0 and  $R^{13}$  is fluoromethyl.

In an embodiment,  $R^{ox}$  is absent.

15 Included in the instant invention is the free form of compounds of Formula I, as well as the pharmaceutically acceptable salts and stereoisomers thereof. Some of the specific compounds exemplified herein are the protonated salts of amine compounds. The term "free form" refers to the amine compounds in non-salt form. The encompassed pharmaceutically acceptable salts not only include the salts exemplified for the specific compounds described herein, but also all the typical pharmaceutically acceptable salts of the free form of compounds of

20 Formula I. The free form of the specific salt compounds described may be isolated using techniques known in the art. For example, the free form may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia and sodium bicarbonate. The free forms may differ from their respective salt forms



somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise pharmaceutically equivalent to their respective free forms for purposes of the invention.

5 The pharmaceutically acceptable salts of the instant compounds can be synthesized from the compounds of this invention which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts of the basic compounds are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents. Similarly, the salts of the acidic compounds are formed by  
10 reactions with the appropriate inorganic or organic base.

Thus, pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed by reacting a basic instant compound with an inorganic or organic acid. For example, conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic,  
15 sulfuric, sulfamic, phosphoric, nitric and the like, as well as salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

20 When the compound of the present invention is acidic, suitable "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc and the like. Particularly preferred are the ammonium,  
25 calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N<sup>1</sup>-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine,  
30 N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

The preparation of the pharmaceutically acceptable salts described above and other typical pharmaceutically acceptable salts is more fully described by Berg *et al.*, "Pharmaceutical Salts," *J. Pharm. Sci.*, 1977:66:1-19.

It will also be noted that the compounds of the present invention are potentially internal salts or zwitterions, since under physiological conditions a deprotonated acidic moiety in the compound, such as a carboxyl group, may be anionic, and this electronic charge might then be balanced off internally against the cationic charge of a protonated or alkylated basic moiety, such as a quaternary nitrogen atom.

The following abbreviations, used in the Schemes and Examples, are defined below:

CDI	1,1'-carbonyldiimidazole
CSP HPLC	Chiral stationary phase high performance liquid chromatography
DAST	(diethylamino)sulfur trifluoride
DCE	1,2-dichloroethane
DCM	Dichloromethane
DMF	Dimethylformamide
DMSO	Dimethyl sulfoxide
EtOAc	Ethyl acetate
LAH	Lithium aluminum hydride
LiHMDS	Lithium hexamethyldisilazide
MsCl	Methanesulfonylchloride
NaHMDS	Sodium hexamethyldisilazide
NOE	Nuclear Overhauser Effect
PTC	Phase transfer catalyst
TBSCl	<i>tert</i> -butyldimethylsilyl chloride
TEA	Triethylamine
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran

The compounds of this invention may be prepared by employing reactions as shown in the following schemes, in addition to other standard manipulations that are known in the literature or exemplified in the experimental procedures. The illustrative schemes below, therefore, are not limited by the compounds listed or by any particular substituents employed for illustrative purposes. Substituent numbering as shown in the schemes does not necessarily correlate to that used in the claims and often, for clarity, a single substituent is shown attached to the compound where multiple substituents are allowed under the definitions of Formula I hereinabove.

## SCHEMES

As shown in Scheme A, key 2,2-disubstituted dihydropyrrole intermediate A-8 may be obtained from readily available suitably substituted  $\alpha$ -phenylglycines. Following the procedure described by Van Betsbrugge et. al. (*Tetrahedron*, 1997, 53, 9233-9240) the  $\alpha$ -allyl- $\alpha$ -phenylglycine A-3 is prepared. Reduction of the ester and cyclization with carbonyldiimidazole provides intermediate A-4. Ruthenium oxidation of the allylic olefin, followed by ester formation and alkylation of the nitrogen provides intermediate A-5. Cyclization and decarboxylation results in intermediate A-6. The ring carbonyl can then be utilized to incorporate a suitably substituted phenyl moiety. Subsequent saponification and oxygen protection leads to the protected intermediate A-8. The enantiomers of A-8 may often be separated utilizing chiral chromatographic techniques. The ring nitrogen may then be reacted with triphosgene to prepare the activated carbonyl chloride A-9.

Scheme B illustrates the preparation of the fluorinated aminopiperidine B-5, starting with the N-protected piperidone. The *cis* and *trans* diastereomeric pairs may often be separated by silica gel chromatography and the enantiomers may often be separated utilizing chiral chromatographic techniques.

As shown in Scheme C, such a fluorinated aminopiperidine may then be reacted with the dihydropyrrole intermediate A-9 to provide the instant compound C-1.

Scheme D illustrates preparation of 2-fluoromethyl-4-aminopiperidine compounds and incorporation of those groups into the compounds of the instant invention. It should be noted that fluoride displacement of the sidechain hydroxyl in intermediate D-3 often leads to both the desired intermediate D-4 and the ring homologous compound D-5. These intermediate compounds may be separated by silica gel chromatography.

Scheme E illustrates an alternative preparation of the 2-fluoromethyl-3-aminopiperidine compounds in which the seven-membered ring isomer is not produced.

As shown in Schemes F and G, the hydroxyl moiety of A-9 may undergo alkylation with a variety of reagents.

As illustrated in Scheme H, replacement of the hydroxyl moiety of compound C-1 with a suitably substituted amine proceeds through the corresponding aldehyde H-1, followed by reductive alkylation, resulting in the instant compound H-2.

Scheme I illustrates the synthesis of 2-monosubstituted dihydropyrrole compounds of the instant invention. Preparation and displacement of the methylimidazolyl moiety in intermediate I-7 with the aminopiperidine provides the instant compound I-8.

Scheme J illustrates homologation of the 2-alkyl sidechain to provide the instant compounds J-3 and J-4.


5 Schemes K to M illustrate further modifications of the C-2 alkyl sidechain starting with the intermediate aldehyde H-1. Thus in Scheme K, the aldehyde H-1 is treated with a Grignard reagent, such as an alkyl Grignard, to provide the hydroxy compound K-1.

10 Scheme L illustrates homologation of the C-2 side chain. The aldehyde H-1 is treated with a phosphonoacetate and the conjugated double bond is then reduced to provide the ester L-1. Subsequent reduction of the ester and oxidation of the alcohol provides the aldehyde L-3, which can reductively alkylate a suitably substituted amine to provide the instant compound L-4. Further alkylation of L-4 is also illustrated.




Scheme M illustrates fluorination of the C-2 sidechain and subsequent conversion of the hydroxyl moiety to an amine via displacement of the corresponding triflate with sodium azide.


15 Scheme N illustrates incorporation of a difluoromethyl moiety into the C-2 sidechain.


## SCHEME A



## SCHEME A (continued)







SCHEME B


SCHEME C




## SCHEME D




SCHEME E

SCHEME F


## SCHEME G






## SCHEME H



## SCHEME I





SCHEME I (continued)


SCHEME J




## SCHEME J (continued)




SCHEME K


SCHEME L

## SCHEME L (continued)



## SCHEME M



SCHEME N

### Utilities

The compounds of the invention find use in a variety of applications. As will be appreciated by those skilled in the art, mitosis may be altered in a variety of ways; that is, one can affect mitosis either by increasing or decreasing the activity of a component in the mitotic pathway. Stated differently, mitosis may be affected (e.g., disrupted) by disturbing equilibrium, either by inhibiting or activating certain components. Similar approaches may be used to alter meiosis.

In a preferred embodiment, the compounds of the invention are used to modulate mitotic spindle formation, thus causing prolonged cell cycle arrest in mitosis. By "modulate" herein is meant altering mitotic spindle formation, including increasing and decreasing spindle formation. By "mitotic spindle formation" herein is meant organization of microtubules into bipolar structures by mitotic kinesins. By "mitotic spindle dysfunction" herein is meant mitotic arrest and monopolar spindle formation.

The compounds of the invention are useful to bind to and/or modulate the activity of a mitotic kinesin. In a preferred embodiment, the mitotic kinesin is a member of the bimC subfamily of mitotic kinesins (as described in U.S. Patent No. 6,284,480, column 5). In a further preferred embodiment, the mitotic kinesin is human KSP, although the activity of mitotic kinesins from other organisms may also be modulated by the compounds of the present invention. In this context, modulate means either increasing or decreasing spindle pole separation, causing malformation, i.e., splaying, of mitotic spindle poles, or otherwise causing morphological perturbation of the mitotic spindle. Also included within the definition of KSP for these purposes are variants and/or fragments of KSP. In addition, other mitotic kinesins may be inhibited by the compounds of the present invention.

The compounds of the invention are used to treat cellular proliferation diseases. Disease states which can be treated by the methods and compositions provided herein include, but are not limited to, cancer (further discussed below), autoimmune disease, arthritis, graft rejection, inflammatory bowel disease, proliferation induced after medical procedures, including, but not limited to, surgery, angioplasty, and the like. It is appreciated that in some cases the cells may not be in a hyper- or hypoproliferation state (abnormal state) and still require treatment. For example, during wound healing, the cells may be proliferating "normally", but proliferation enhancement may be desired. Similarly, as discussed above, in the agriculture arena, cells may be in a "normal" state, but proliferation modulation may be desired to enhance a crop by directly enhancing growth of a crop, or by inhibiting the growth of a plant or organism which adversely

affects the crop. Thus, in one embodiment, the invention herein includes application to cells or individuals afflicted or impending affliction with any one of these disorders or states.

The compounds, compositions and methods provided herein are particularly deemed useful for the treatment of cancer including solid tumors such as skin, breast, brain, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that may be treated by the compounds, compositions and methods of the invention include, but are not limited to:

Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous



cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin:  
5 malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles, dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above-identified conditions.

10 The compounds of the instant invention may also be useful as antifungal agents, by modulating the activity of the fungal members of the bimC kinesin subgroup, as is described in U.S. Patent No. 6,284,480.

The compounds of this invention may be administered to mammals, preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers, excipients or diluents, in a pharmaceutical composition, according to standard pharmaceutical  
15 practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specific amounts, as well as any product which results, directly or indirectly, from combination of the specific ingredients in the specified  
20 amounts.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art  
25 for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be  
30 for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, microcrystalline cellulose, sodium crosscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by  
35 known techniques to mask the unpleasant taste of the drug or delay disintegration and absorption

in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water soluble taste masking material such as hydroxypropyl-methylcellulose or hydroxypropylcellulose, or a time delay material such as ethyl cellulose, cellulose acetate butyrate may be employed.

5                   Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

10                   Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products  
15 of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol  
20 anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

25                   Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated hydroxyanisole or alpha-tocopherol.

30                   Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. These  
35 compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally occurring phosphatides, for example soy bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavoring agents, preservatives and antioxidants.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, flavoring and coloring agents and antioxidant.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous solutions. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

The sterile injectable preparation may also be a sterile injectable oil-in-water microemulsion where the active ingredient is dissolved in the oily phase. For example, the active ingredient may be first dissolved in a mixture of soybean oil and lecithin. The oil solution then introduced into a water and glycerol mixture and processed to form a microemulsion.

The injectable solutions or microemulsions may be introduced into a patient's blood stream by local bolus injection. Alternatively, it may be advantageous to administer the solution or microemulsion in such a way as to maintain a constant circulating concentration of the instant compound. In order to maintain such a constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension for intramuscular and subcutaneous administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butane diol. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Compounds of Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug

with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

5 For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compound of Formula I are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or via transdermal routes,  
10 using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Compounds of the present invention may also be delivered as a suppository employing bases such as cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of  
15 various molecular weights and fatty acid esters of polyethylene glycol.

When a compound according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, sex and response of the individual patient, as well as the severity of the patient's symptoms.

20 In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day, preferably of between 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day.

The instant compounds are also useful in combination with known therapeutic  
25 agents and anti-cancer agents. For example, instant compounds are useful in combination with known anti-cancer agents. Combinations of the presently disclosed compounds with other anti-cancer or chemotherapeutic agents are within the scope of the invention. Examples of such agents can be found in *Cancer Principles and Practice of Oncology* by V.T. Devita and S. Hellman (editors), 6<sup>th</sup> edition (February 15, 2001), Lippincott Williams & Wilkins Publishers. A  
30 person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the cancer involved. Such anti-cancer agents include, but are not limited to, the following: estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic/cytostatic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors and  
35 other angiogenesis inhibitors, inhibitors of cell proliferation and survival signaling, apoptosis

inducing agents and agents that interfere with cell cycle checkpoints. The instant compounds are particularly useful when co-administered with radiation therapy.

In an embodiment, the instant compounds are also useful in combination with known anti-cancer agents including the following: estrogen receptor modulators, androgen  
5 receptor modulators, retinoid receptor modulators, cytotoxic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors, and other angiogenesis inhibitors.

“Estrogen receptor modulators” refers to compounds that interfere with or inhibit the binding of estrogen to the receptor, regardless of mechanism. Examples of estrogen receptor  
10 modulators include, but are not limited to, tamoxifen, raloxifene, idoxifene, LY353381, LY117081, toremifene, fulvestrant, 4-[7-(2,2-dimethyl-1-oxopropoxy-4-methyl-2-[4-[2-(1-piperidinyl)ethoxy]phenyl]-2H-1-benzopyran-3-yl)]-phenyl-2,2-dimethylpropanoate, 4,4'-dihydroxybenzophenone-2,4-dinitrophenyl-hydrazone, and SH646.

“Androgen receptor modulators” refers to compounds which interfere or inhibit  
15 the binding of androgens to the receptor, regardless of mechanism. Examples of androgen receptor modulators include finasteride and other 5 $\alpha$ -reductase inhibitors, nilutamide, flutamide, bicalutamide, liarozole, and abiraterone acetate.

“Retinoid receptor modulators” refers to compounds which interfere or inhibit the binding of retinoids to the receptor, regardless of mechanism. Examples of such retinoid  
20 receptor modulators include bexarotene, tretinoin, 13-cis-retinoic acid, 9-cis-retinoic acid,  $\alpha$ -difluoromethylornithine, ILX23-7553, trans-N-(4'-hydroxyphenyl) retinamide, and N-4-carboxyphenyl retinamide.

“Cytotoxic/cytostatic agents” refer to compounds which cause cell death or inhibit cell proliferation primarily by interfering directly with the cell's functioning or inhibit or  
25 interfere with cell mytosis, including alkylating agents, tumor necrosis factors, intercalators, hypoxia activatable compounds, microtubule inhibitors/microtubule-stabilizing agents, inhibitors of mitotic kinesins, inhibitors of kinases involved in mitotic progression, antimetabolites; biological response modifiers; hormonal/anti-hormonal therapeutic agents, haematopoietic growth factors, monoclonal antibody targeted therapeutic agents, topoisomerase inhibitors,  
30 proteasome inhibitors and ubiquitin ligase inhibitors.

Examples of cytotoxic agents include, but are not limited to, sertenef, cachectin, ifosfamide, tasonermin, lonidamine, carboplatin, altretamine, prednimustine, dibromodulcitol, ranimustine, fotemustine, nedaplatin, oxaliplatin, temozolomide, heptaplatin, estramustine, improsulfan tosilate, trofosfamide, nimustine, dibrospidium chloride, pumitepa, lobaplatin,  
35 satraplatin, proflomycin, cisplatin, irofulven, dexifosfamide, cis-aminedichloro(2-methyl-

pyridine)platinum, benzylguanine, glufosfamide, GPX100, (trans, trans, trans)-bis-mu-(hexane-1,6-diamine)-mu-[diamine-platinum(II)]bis[diamine(chloro)platinum (II)]tetrachloride, diarizidinylspermine, arsenic trioxide, 1-(11-dodecylamino-10-hydroxyundecyl)-3,7-dimethylxanthine, zorubicin, idarubicin, daunorubicin, bisantrene, mitoxantrone, pirarubicin, pinafide, valrubicin, amrubicin, antineoplaston, 3'-deamino-3'-morpholino-13-deoxo-10-hydroxycarminomycin, annamycin, galarubicin, elinafide, MEN10755, and 4-demethoxy-3-deamino-3-aziridinyl-4-methylsulphonyl-daunorubicin (see WO 00/50032).

An example of a hypoxia activatable compound is tirapazamine.

Examples of proteasome inhibitors include but are not limited to lactacystin and

10 bortezomib

Examples of microtubule inhibitors/microtubule-stabilising agents include paclitaxel, vindesine sulfate, 3',4'-didehydro-4'-deoxy-8'-norvincal leukoblastine, docetaxol, rhizoxin, dolastatin, mivobulin isethionate, auristatin, cemadotin, RPR109881, BMS184476, vinflunine, cryptophycin, 2,3,4,5,6-pentafluoro-N-(3-fluoro-4-methoxyphenyl) benzene sulfonamide, anhydrovinblastine, N,N-dimethyl-L-valyl-L-valyl-N-methyl-L-valyl-L-prolyl-L-proline-t-butylamide, TDX258, the epothilones (see for example U.S. Pat. Nos. 6,284,781 and 6,288,237) and BMS188797.

Some examples of topoisomerase inhibitors are topotecan, hycaptamine, irinotecan, rubitecan, 6-ethoxypropionyl-3',4'-O-exo-benzylidene-chartreusin, 9-methoxy-N,N-dimethyl-5-nitropyrazolo[3,4,5-kl]acridine-2-(6H) propanamine, 1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4':b,7]-indolizino[1,2b]quinoline-10,13(9H,15H)dione, lurtotecan, 7-[2-(N-isopropylamino)ethyl]-(20S)camptothecin, BNP1350, BNPI1100, BN80915, BN80942, etoposide phosphate, teniposide, sobuzoxane, 2'-dimethylamino-2'-deoxy-etoposide, GL331, N-[2-(dimethylamino)ethyl]-9-hydroxy-5,6-dimethyl-6H-pyrido[4,3-b]carbazole-1-carboxamide, asulacrine, (5a, 5aB, 8aa,9b)-9-[2-[N-[2-(dimethylamino)ethyl]-N-methylamino]ethyl]-5-[4-hydroxy-3,5-dimethoxyphenyl]-5,5a,6,8,8a,9-hexahydrofuro(3',4':6,7)naphtho(2,3-d)-1,3-dioxol-6-one, 2,3-(methylenedioxy)-5-methyl-7-hydroxy-8-methoxybenzo[c]-phenanthridinium, 6,9-bis[(2-aminoethyl)amino]benzo[g]isoguinoline-5,10-dione, 5-(3-aminopropylamino)-7,10-dihydroxy-2-(2-hydroxyethylaminomethyl)-6H-pyrazolo[4,5,1-de]acridin-6-one, N-[1-[2(diethylamino)ethylamino]-7-methoxy-9-oxo-9H-thioxanthen-4-ylmethyl]formamide, N-(2-(dimethylamino)ethyl)acridine-4-carboxamide, 6-[[2-(dimethylamino)ethyl]amino]-3-hydroxy-7H-indeno[2,1-c] quinolin-7-one, and dimesna.

Examples of inhibitors of mitotic kinesins, and in particular the human mitotic kinesin KSP, are described in PCT Publications WO 01/30768 and WO 01/98278, WO

03/050,064 (June 19, 2003), WO 03/050,122 (June 19, 2003), WO 03/049,527 (June 19, 2003), WO 03/049,679 (June 19, 2003), WO 03/049,678 (June 19, 2003) and WO 03/39460 (May 15, 2003) and pending PCT Appl. Nos. US03/06403 (filed March 4, 2003), US03/15861 (filed May 19, 2003), US03/15810 (filed May 19, 2003), US03/18482 (filed June 12, 2003) and

5 US03/18694 (filed June 12, 2003). In an embodiment inhibitors of mitotic kinesins include, but are not limited to inhibitors of KSP, inhibitors of MKLP1, inhibitors of CENP-E, inhibitors of MCAK, inhibitors of Kif14, inhibitors of Mphosph1 and inhibitors of Rab6-KIFL.

“Inhibitors of kinases involved in mitotic progression” include, but are not limited to, inhibitors of aurora kinase, inhibitors of Polo-like kinases (PLK) (in particular inhibitors of  
10 PLK-1), inhibitors of bub-1 and inhibitors of bub-R1.


“Antiproliferative agents” includes antisense RNA and DNA oligonucleotides such as G3139, ODN698, RVASKRAS, GEM231, and INX3001, and antimetabolites such as enocitabine, carmofur, tegafur, pentostatin, doxifluridine, trimetrexate, fludarabine, capecitabine, galocitabine, cytarabine ocfosfate, fosteabine sodium hydrate, raltitrexed, paltitrexid, emitefur,  
15 tiazofurin, decitabine, nolatrexed, pemetrexed, nelzarabine, 2'-deoxy-2'-methylidenecytidine, 2'-fluoromethylene-2'-deoxycytidine, N-[5-(2,3-dihydro-benzofuryl)sulfonyl]-N'-(3,4-dichlorophenyl)urea, N6-[4-deoxy-4-[N2-[2(E),4(E)-tetradecadienoyl]glycylamino]-L-glycero-B-L-manno-heptopyranosyl]adenine, aplidine, ecteinascidin, troxacitabine, 4-[2-amino-4-oxo-4,6,7,8-tetrahydro-3H-pyrimidino[5,4-b][1,4]thiazin-6-yl-(S)-ethyl]-2,5-thienoyl-L-glutamic  
20 acid, aminopterin, 5-fluorouracil, alanosine, 11-acetyl-8-(carbamoxyloxymethyl)-4-formyl-6-methoxy-14-oxa-1,11-diazatetracyclo(7.4.1.0.0)-tetradeca-2,4,6-trien-9-yl acetic acid ester, swainsonine, lometrexol, dexrazoxane, methioninase, 2'-cyano-2'-deoxy-N4-palmitoyl-1-B-D-arabino furanosyl cytosine and 3-aminopyridine-2-carboxaldehyde thiosemicarbazone.

Examples of monoclonal antibody targeted therapeutic agents include those  
25 therapeutic agents which have cytotoxic agents or radioisotopes attached to a cancer cell specific or target cell specific monoclonal antibody. Examples include Bexxar.

“HMG-CoA reductase inhibitors” refers to inhibitors of 3-hydroxy-3-methylglutaryl-CoA reductase. Compounds which have inhibitory activity for HMG-CoA reductase can be readily identified by using assays well-known in the art. For example, see the  
30 assays described or cited in U.S. Patent 4,231,938 at col. 6, and WO 84/02131 at pp. 30-33. The terms “HMG-CoA reductase inhibitor” and “inhibitor of HMG-CoA reductase” have the same meaning when used herein.

Examples of HMG-CoA reductase inhibitors that may be used include but are not limited to lovastatin (MEVACOR®; see U.S. Patent Nos. 4,231,938, 4,294,926 and 4,319,039),  
35 simvastatin (ZOCOR®; see U.S. Patent Nos. 4,444,784, 4,820,850 and 4,916,239), pravastatin

(PRAVACHOL®; see U.S. Patent Nos. 4,346,227, 4,537,859, 4,410,629, 5,030,447 and 5,180,589), fluvastatin (LESCOL®; see U.S. Patent Nos. 5,354,772, 4,911,165, 4,929,437, 5,189,164, 5,118,853, 5,290,946 and 5,356,896) and atorvastatin (LIPITOR®; see U.S. Patent Nos. 5,273,995, 4,681,893, 5,489,691 and 5,342,952). The structural formulas of these and additional HMG-CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", *Chemistry & Industry*, pp. 85-89 (5 February 1996) and US Patent Nos. 4,782,084 and 4,885,314. The term HMG-CoA reductase inhibitor as used herein includes all pharmaceutically acceptable lactone and open-acid forms (i.e., where the lactone ring is opened to form the free acid) as well as salt and ester forms of compounds which have HMG-CoA reductase inhibitory activity, and therefor the use of such salts, esters, open-acid and lactone forms is included within the scope of this invention. An illustration of the lactone portion and its corresponding open-acid form is shown below as structures I and II.



In HMG-CoA reductase inhibitors where an open-acid form can exist, salt and ester forms may be formed from the open-acid, and all such forms are included within the meaning of the term "HMG-CoA reductase inhibitor" as used herein. In an embodiment, the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin, and in a further embodiment, simvastatin. Herein, the term "pharmaceutically acceptable salts" with respect to the HMG-CoA reductase inhibitor shall mean non-toxic salts of the compounds employed in this invention which are generally prepared by reacting the free acid with a suitable organic or inorganic base, particularly those formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc and tetramethylammonium, as well as those salts formed from amines such as ammonia, ethylenediamine, N-methylglucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, 1-p-chlorobenzyl-2-pyrrolidine-1'-yl-methylbenz-imidazole, diethylamine, piperazine, and tris(hydroxymethyl) aminomethane. Further examples of salt



forms of HMG-CoA reductase inhibitors may include, but are not limited to, acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, 5 hydrabamine, hydrobromide, hydrochloride, hydroxynapthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamaote, palmitate, panthothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide, and valerate.

10 Ester derivatives of the described HMG-CoA reductase inhibitor compounds may act as prodrugs which, when absorbed into the bloodstream of a warm-blooded animal, may cleave in such a manner as to release the drug form and permit the drug to afford improved therapeutic efficacy.

“Prenyl-protein transferase inhibitor” refers to a compound which inhibits any 15 one or any combination of the prenyl-protein transferase enzymes, including farnesyl-protein transferase (FPTase), geranylgeranyl-protein transferase type I (GGPTase-I), and geranylgeranyl-protein transferase type-II (GGPTase-II, also called Rab GGPTase). Examples of prenyl-protein transferase inhibiting compounds include (+)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, (-)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, (+)-6- 20 [amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl) methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, 5(S)-n-butyl-1-(2,3-dimethylphenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone, (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl) methyl]-2-piperazinone, 5(S)-n-Butyl-1-(2-methylphenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone, 1-(3-chlorophenyl)- 4-[1-(4-cyanobenzyl)-2-methyl-5-imidazolylmethyl]-2-piperazinone, 1-(2,2-diphenylethyl)-3-[N-(1-(4-cyanobenzyl)-1H-imidazol-5-ylethyl)carbamoyl]piperidine, 4-{5-[4-hydroxymethyl-4-(4-chloropyridin-2-ylmethyl)-piperidine-1-ylmethyl]-2-methylimidazol-1-ylmethyl} benzonitrile, 4- 25 {5-[4-hydroxymethyl-4-(3-chlorobenzyl)-piperidine-1-ylmethyl]-2-methylimidazol-1-ylmethyl} benzonitrile, 4-{3-[4-(2-oxo-2H-pyridin-1-yl)benzyl]-3H-imidazol-4-ylmethyl} benzonitrile, 4-{3-[4-(5-chloro-2-oxo-2H-[1,2']bipyridin-5'-ylmethyl)-3H-imidazol-4-ylmethyl} benzonitrile, 4-{3-[4-(2-oxo-2H-[1,2'] bipyridin-5'-ylmethyl)-3H-imidazol-4-ylmethyl} benzonitrile, 4-[3-(2-oxo-1-phenyl-1,2-dihydropyridin-4-ylmethyl)-3H-imidazol-4-ylmethyl] benzonitrile, 18,19-dihydro-19-oxo-5H,17H-6,10:12,16-dimetheno-1H-imidazo[4,3- 35 c][1,11,4]dioxazacyclo-nonadecine-9-carbonitrile, (±)-19,20-dihydro-19-oxo-5H-18,21-ethano-

12,14-etheno-6,10-metheno-22*H*-benzo[*d*]imidazo[4,3-*k*][1,6,9,12]oxatriaza-cyclooctadecine-9-carbonitrile, 19,20-dihydro-19-oxo-5*H*,17*H*-18,21-ethano-6,10:12,16-dimetheno-22*H*-imidazo[3,4-*h*][1,8,11,14]oxatriazacycloeicosine-9-carbonitrile, and (±)-19,20-dihydro-3-methyl-19-oxo-5*H*-18,21-ethano-12,14-etheno-6,10-metheno-22*H*-benzo[*d*]imidazo[4,3-*k*][1,6,9,12]oxa-triazacyclooctadecine-9-carbonitrile.

Other examples of prenyl-protein transferase inhibitors can be found in the following publications and patents: WO 96/30343, WO 97/18813, WO 97/21701, WO 97/23478, WO 97/38665, WO 98/28980, WO 98/29119, WO 95/32987, U.S. Patent No. 5,420,245, U.S. Patent No. 5,523,430, U.S. Patent No. 5,532,359, U.S. Patent No. 5,510,510, U.S. Patent No. 5,589,485, U.S. Patent No. 5,602,098, European Patent Publ. 0 618 221, European Patent Publ. 0 675 112, European Patent Publ. 0 604 181, European Patent Publ. 0 696 593, WO 94/19357, WO 95/08542, WO 95/11917, WO 95/12612, WO 95/12572, WO 95/10514, U.S. Patent No. 5,661,152, WO 95/10515, WO 95/10516, WO 95/24612, WO 95/34535, WO 95/25086, WO 96/05529, WO 96/06138, WO 96/06193, WO 96/16443, WO 96/21701, WO 96/21456, WO 96/22278, WO 96/24611, WO 96/24612, WO 96/05168, WO 96/05169, WO 96/00736, U.S. Patent No. 5,571,792, WO 96/17861, WO 96/33159, WO 96/34850, WO 96/34851, WO 96/30017, WO 96/30018, WO 96/30362, WO 96/30363, WO 96/31111, WO 96/31477, WO 96/31478, WO 96/31501, WO 97/00252, WO 97/03047, WO 97/03050, WO 97/04785, WO 97/02920, WO 97/17070, WO 97/23478, WO 97/26246, WO 97/30053, WO 97/44350, WO 98/02436, and U.S. Patent No. 5,532,359.

For an example of the role of a prenyl-protein transferase inhibitor on angiogenesis see European J. of Cancer, Vol. 35, No. 9, pp.1394-1401 (1999).

"Angiogenesis inhibitors" refers to compounds that inhibit the formation of new blood vessels, regardless of mechanism. Examples of angiogenesis inhibitors include, but are not limited to, tyrosine kinase inhibitors, such as inhibitors of the tyrosine kinase receptors Flt-1 (VEGFR1) and Flk-1/KDR (VEGFR2), inhibitors of epidermal-derived, fibroblast-derived, or platelet derived growth factors, MMP (matrix metalloprotease) inhibitors, integrin blockers, interferon- $\alpha$ , interleukin-12, pentosan polysulfate, cyclooxygenase inhibitors, including nonsteroidal anti-inflammatories (NSAIDs) like aspirin and ibuprofen as well as selective cyclooxygenase-2 inhibitors like celecoxib and rofecoxib (PNAS, Vol. 89, p. 7384 (1992); JNCI, Vol. 69, p. 475 (1982); Arch. Ophthalmol., Vol. 108, p.573 (1990); Anat. Rec., Vol. 238, p. 68 (1994); FEBS Letters, Vol. 372, p. 83 (1995); Clin. Orthop. Vol. 313, p. 76 (1995); J. Mol. Endocrinol., Vol. 16, p.107 (1996); Jpn. J. Pharmacol., Vol. 75, p. 105 (1997); Cancer Res., Vol. 57, p. 1625 (1997); Cell, Vol. 93, p. 705 (1998); Intl. J. Mol. Med., Vol. 2, p. 715 (1998); J. Biol. Chem., Vol. 274, p. 9116 (1999)), steroidal anti-inflammatories (such as corticosteroids,

mineralocorticoids, dexamethasone, prednisone, prednisolone, methylpred, betamethasone), carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, angiotensin II antagonists (see Fernandez et al., J. Lab. Clin. Med. 105:141-145 (1985)), and antibodies to VEGF (see, Nature Biotechnology, Vol. 17, pp.963-968 (October 1999); Kim et al., Nature, 362, 841-844 (1993); WO 00/44777; and WO 00/61186).

Other therapeutic agents that modulate or inhibit angiogenesis and may also be used in combination with the compounds of the instant invention include agents that modulate or inhibit the coagulation and fibrinolysis systems (see review in *Clin. Chem. La. Med.* 38:679-692 (2000)). Examples of such agents that modulate or inhibit the coagulation and fibrinolysis pathways include, but are not limited to, heparin (see *Thromb. Haemost.* 80:10-23 (1998)), low molecular weight heparins and carboxypeptidase U inhibitors (also known as inhibitors of active thrombin activatable fibrinolysis inhibitor [TAFIa]) (see *Thrombosis Res.* 101:329-354 (2001)). TAFIa inhibitors have been described in PCT Publication WO 03/013,526 and U.S. Ser. No. 60/349,925 (filed January 18, 2002).

“Agents that interfere with cell cycle checkpoints” refer to compounds that inhibit protein kinases that transduce cell cycle checkpoint signals, thereby sensitizing the cancer cell to DNA damaging agents. Such agents include inhibitors of ATR, ATM, the Chk1 and Chk2 kinases and cdk and cdc kinase inhibitors and are specifically exemplified by 7-hydroxystaurosporin, flavopiridol, CYC202 (Cyclacel) and BMS-387032.

“Inhibitors of cell proliferation and survival signaling pathway” refer to pharmaceutical agents that inhibit cell surface receptors and signal transduction cascades downstream of those surface receptors. Such agents include inhibitors of inhibitors of EGFR (for example gefitinib and erlotinib), inhibitors of ERB-2 (for example trastuzumab), inhibitors of IGFR, inhibitors of cytokine receptors, inhibitors of MET, inhibitors of PI3K (for example LY294002), serine/threonine kinases (including but not limited to inhibitors of Akt such as described in WO 02/083064, WO 02/083139, WO 02/083140 and WO 02/083138), inhibitors of Raf kinase (for example BAY-43-9006), inhibitors of MEK (for example CI-1040 and PD-098059) and inhibitors of mTOR (for example Wyeth CCI-779). Such agents include small molecule inhibitor compounds and antibody antagonists.

“Apoptosis inducing agents” include activators of TNF receptor family members (including the TRAIL receptors).


The combinations with NSAID's are directed to the use of NSAID's which are potent COX-2 inhibiting agents. For purposes of this specification an NSAID is potent if it

possesses an  $IC_{50}$  for the inhibition of COX-2 of  $1\mu M$  or less as measured by cell or microsomal assays.


The invention also encompasses combinations with NSAID's which are selective COX-2 inhibitors. For purposes of this specification NSAID's which are selective inhibitors of COX-2 are defined as those which possess a specificity for inhibiting COX-2 over COX-1 of at least 100 fold as measured by the ratio of  $IC_{50}$  for COX-2 over  $IC_{50}$  for COX-1 evaluated by cell or microsomal assays. Such compounds include, but are not limited to those disclosed in U.S. Patent 5,474,995, issued December 12, 1995, U.S. Patent 5,861,419, issued January 19, 1999, U.S. Patent 6,001,843, issued December 14, 1999, U.S. Patent 6,020,343, issued February 1, 2000, U.S. Patent 5,409,944, issued April 25, 1995, U.S. Patent 5,436,265, issued July 25, 1995, U.S. Patent 5,536,752, issued July 16, 1996, U.S. Patent 5,550,142, issued August 27, 1996, U.S. Patent 5,604,260, issued February 18, 1997, U.S. 5,698,584, issued December 16, 1997, U.S. Patent 5,710,140, issued January 20, 1998, WO 94/15932, published July 21, 1994, U.S. Patent 5,344,991, issued June 6, 1994, U.S. Patent 5,134,142, issued July 28, 1992, U.S. Patent 5,380,738, issued January 10, 1995, U.S. Patent 5,393,790, issued February 20, 1995, U.S. Patent 5,466,823, issued November 14, 1995, U.S. Patent 5,633,272, issued May 27, 1997, and U.S. Patent 5,932,598, issued August 3, 1999, all of which are hereby incorporated by reference.

Inhibitors of COX-2 that are particularly useful in the instant method of treatment are:

3-phenyl-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone; and





5-chloro-3-(4-(methylsulfonyl)phenyl)-2-(2-methyl-5-pyridinyl)pyridine;




or a pharmaceutically acceptable salt thereof.

General and specific synthetic procedures for the preparation of the COX-2 inhibitor compounds described above are found in U.S. Patent No. 5,474,995, issued December 12, 1995, U.S. Patent No. 5,861,419, issued January 19, 1999, and U.S. Patent No. 6,001,843, issued December 14, 1999, all of which are herein incorporated by reference.

Compounds that have been described as specific inhibitors of COX-2 and are therefore useful in the present invention include, but are not limited to, the following:





or a pharmaceutically acceptable salt thereof.

Compounds which are described as specific inhibitors of COX-2 and are therefore useful in the present invention, and methods of synthesis thereof, can be found in the following patents, pending applications and publications, which are herein incorporated by reference: WO 94/15932, published July 21, 1994, U.S. Patent No. 5,344,991, issued June 6, 1994, U.S. Patent No. 5,134,142, issued July 28, 1992, U.S. Patent No. 5,380,738, issued January 10, 1995, U.S. Patent No. 5,393,790, issued February 20, 1995, U.S. Patent No. 5,466,823, issued November 14, 1995, U.S. Patent No. 5,633,272, issued May 27, 1997, and U.S. Patent No. 5,932,598, issued August 3, 1999.

Compounds which are specific inhibitors of COX-2 and are therefore useful in the present invention, and methods of synthesis thereof, can be found in the following patents, pending applications and publications, which are herein incorporated by reference: U.S. Patent No. 5,474,995, issued December 12, 1995, U.S. Patent No. 5,861,419, issued January 19, 1999, U.S. Patent No. 6,001,843, issued December 14, 1999, U.S. Patent No. 6,020,343, issued February 1, 2000, U.S. Patent No. 5,409,944, issued April 25, 1995, U.S. Patent No. 5,436,265, issued July 25, 1995, U.S. Patent No. 5,536,752, issued July 16, 1996, U.S. Patent No. 5,550,142, issued August 27, 1996, U.S. Patent No. 5,604,260, issued February 18, 1997, U.S. Patent No. 5,698,584, issued December 16, 1997, and U.S. Patent No. 5,710,140, issued January 20, 1998.

Other examples of angiogenesis inhibitors include, but are not limited to, endostatin, ukrain, ranpirase, IM862, 5-methoxy-4-[2-methyl-3-(3-methyl-2-butenyl)oxiranyl]-1-oxaspiro[2,5]oct-6-yl(chloroacetyl)carbamate, acetyldinanaline, 5-amino-1-[[3,5-dichloro-4-(4-chlorobenzoyl)phenyl]methyl]-1H-1,2,3-triazole-4-carboxamide, CM101, squalamine, combretastatin, RPI4610, NX31838, sulfated mannopentaose phosphate, 7,7-(carbonyl-bis[imino-N-methyl-4,2-pyrrolocarbonylimino[N-methyl-4,2-pyrrole]-carbonylimino]-bis-(1,3-naphthalene disulfonate), and 3-[(2,4-dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416).

As used above, "integrin blockers" refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the  $\alpha_v\beta_3$  integrin, to

compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the  $\alpha_v\beta_5$  integrin, to compounds which antagonize, inhibit or counteract binding of a physiological ligand to both the  $\alpha_v\beta_3$  integrin and the  $\alpha_v\beta_5$  integrin, and to compounds which antagonize, inhibit or counteract the activity of the particular integrin(s) expressed on capillary endothelial cells. The term also refers to antagonists of the  $\alpha_v\beta_6$ ,  $\alpha_v\beta_8$ ,  $\alpha_1\beta_1$ ,  $\alpha_2\beta_1$ ,  $\alpha_5\beta_1$ ,  $\alpha_6\beta_1$  and  $\alpha_6\beta_4$  integrins. The term also refers to antagonists of any combination of  $\alpha_v\beta_3$ ,  $\alpha_v\beta_5$ ,  $\alpha_v\beta_6$ ,  $\alpha_v\beta_8$ ,  $\alpha_1\beta_1$ ,  $\alpha_2\beta_1$ ,  $\alpha_5\beta_1$ ,  $\alpha_6\beta_1$  and  $\alpha_6\beta_4$  integrins.

Some specific examples of tyrosine kinase inhibitors include N-(trifluoromethylphenyl)-5-methylisoxazol-4-carboxamide, 3-[(2,4-dimethylpyrrol-5-yl)methylidenyl]indolin-2-one, 17-(allylamino)-17-demethoxygeldanamycin, 4-(3-chloro-4-fluorophenylamino)-7-methoxy-6-[3-(4-morpholinyl)propoxyl]quinazoline, N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, BIBX1382, 2,3,9,10,11,12-hexahydro-10-(hydroxymethyl)-10-hydroxy-9-methyl-9,12-epoxy-1H-diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-i][1,6]benzodiazocin-1-one, SH268, genistein, STI571, CEP2563, 4-(3-chlorophenylamino)-5,6-dimethyl-7H-pyrrolo[2,3-d]pyrimidinemethane sulfonate, 4-(3-bromo-4-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, 4-(4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, SU6668, STI571A, N-4-chlorophenyl-4-(4-pyridylmethyl)-1-phthalazinamine, and EMD121974.

Combinations with compounds other than anti-cancer compounds are also encompassed in the instant methods. For example, combinations of the instantly claimed compounds with PPAR- $\gamma$  (i.e., PPAR-gamma) agonists and PPAR- $\delta$  (i.e., PPAR-delta) agonists are useful in the treatment of certain malignancies. PPAR- $\gamma$  and PPAR- $\delta$  are the nuclear peroxisome proliferator-activated receptors  $\gamma$  and  $\delta$ . The expression of PPAR- $\gamma$  on endothelial cells and its involvement in angiogenesis has been reported in the literature (see *J. Cardiovasc. Pharmacol.* 1998; 31:909-913; *J. Biol. Chem.* 1999; 274:9116-9121; *Invest. Ophthalmol Vis. Sci.* 2000; 41:2309-2317). More recently, PPAR- $\gamma$  agonists have been shown to inhibit the angiogenic response to VEGF in vitro; both troglitazone and rosiglitazone maleate inhibit the development of retinal neovascularization in mice. (*Arch. Ophthalmol.* 2001; 119:709-717). Examples of PPAR- $\gamma$  agonists and PPAR-  $\gamma/\alpha$  agonists include, but are not limited to, thiazolidinediones (such as DRF2725, CS-011, troglitazone, rosiglitazone, and pioglitazone), fenofibrate, gemfibrozil, clofibrate, GW2570, SB219994, AR-H039242, JTT-501, MCC-555, GW2331, GW409544, NN2344, KRP297, NP0110, DRF4158, NN622, GI262570, PNU182716, DRF552926, 2-[(5,7-dipropyl-3-trifluoromethyl-1,2-benzisoxazol-6-yl)oxy]-2-methylpropionic acid (disclosed in USSN 09/782,856), and 2(R)-7-(3-(2-chloro-4-(4-fluorophenoxy)

phenoxy)propoxy)-2-ethylchromane-2-carboxylic acid (disclosed in USSN 60/235,708 and 60/244,697).

Another embodiment of the instant invention is the use of the presently disclosed compounds in combination with gene therapy for the treatment of cancer. For an overview of genetic strategies to treating cancer see Hall et al (*Am J Hum Genet* 61:785-789, 1997) and Kufe et al (*Cancer Medicine*, 5th Ed, pp 876-889, BC Decker, Hamilton 2000). Gene therapy can be used to deliver any tumor suppressing gene. Examples of such genes include, but are not limited to, p53, which can be delivered via recombinant virus-mediated gene transfer (see U.S. Patent No. 6,069,134, for example), a uPA/uPAR antagonist ("Adenovirus-Mediated Delivery of a uPA/uPAR Antagonist Suppresses Angiogenesis-Dependent Tumor Growth and Dissemination in Mice," *Gene Therapy*, August 1998;5(8):1105-13), and interferon gamma (*J Immunol* 2000;164:217-222).

The compounds of the instant invention may also be administered in combination with an inhibitor of inherent multidrug resistance (MDR), in particular MDR associated with high levels of expression of transporter proteins. Such MDR inhibitors include inhibitors of p-glycoprotein (P-gp), such as LY335979, XR9576, OC144-093, R101922, VX853 and PSC833 (valspodar).

A compound of the present invention may be employed in conjunction with anti-emetic agents to treat nausea or emesis, including acute, delayed, late-phase, and anticipatory emesis, which may result from the use of a compound of the present invention, alone or with radiation therapy. For the prevention or treatment of emesis, a compound of the present invention may be used in conjunction with other anti-emetic agents, especially neurokinin-1 receptor antagonists, 5HT3 receptor antagonists, such as ondansetron, granisetron, tropisetron, and zatisetron, GABAB receptor agonists, such as baclofen, a corticosteroid such as Decadron (dexamethasone), Kenalog, Aristocort, Nasalide, Preferid, Benecorten or others such as disclosed in U.S. Patent Nos. 2,789,118, 2,990,401, 3,048,581, 3,126,375, 3,929,768, 3,996,359, 3,928,326 and 3,749,712, an antidopaminergic, such as the phenothiazines (for example prochlorperazine, fluphenazine, thioridazine and mesoridazine), metoclopramide or dronabinol. For the treatment or prevention of emesis that may result upon administration of the instant compounds, conjunctive therapy with an anti-emesis agent selected from a neurokinin-1 receptor antagonist, a 5HT3 receptor antagonist and a corticosteroid is preferred.

Neurokinin-1 receptor antagonists of use in conjunction with the compounds of the present invention are fully described, for example, in U.S. Patent Nos. 5,162,339, 5,232,929, 5,242,930, 5,373,003, 5,387,595, 5,459,270, 5,494,926, 5,496,833, 5,637,699, 5,719,147; European Patent Publication Nos. EP 0 360 390, 0 394 989, 0 428 434, 0 429 366, 0 430 771, 0



436 334, 0 443 132, 0 482 539, 0 498 069, 0 499 313, 0 512 901, 0 512 902, 0 514 273, 0 514 274, 0 514 275, 0 514 276, 0 515 681, 0 517 589, 0 520 555, 0 522 808, 0 528 495, 0 532 456, 0 533 280, 0 536 817, 0 545 478, 0 558 156, 0 577 394, 0 585 913, 0 590 152, 0 599 538, 0 610 793, 0 634 402, 0 686 629, 0 693 489, 0 694 535, 0 699 655, 0 699 674, 0 707 006, 0 708 101, 0 709 375, 0 709 376, 0 714 891, 0 723 959, 0 733 632 and 0 776 893; PCT International Patent Publication Nos. WO 90/05525, 90/05729, 91/09844, 91/18899, 92/01688, 92/06079, 92/12151, 92/15585, 92/17449, 92/20661, 92/20676, 92/21677, 92/22569, 93/00330, 93/00331, 93/01159, 93/01165, 93/01169, 93/01170, 93/06099, 93/09116, 93/10073, 93/14084, 93/14113, 93/18023, 93/19064, 93/21155, 93/21181, 93/23380, 93/24465, 94/00440, 94/01402, 94/02461, 94/02595, 94/03429, 94/03445, 94/04494, 94/04496, 94/05625, 94/07843, 94/08997, 94/10165, 94/10167, 94/10168, 94/10170, 94/11368, 94/13639, 94/13663, 94/14767, 94/15903, 94/19320, 94/19323, 94/20500, 94/26735, 94/26740, 94/29309, 95/02595, 95/04040, 95/04042, 95/06645, 95/07886, 95/07908, 95/08549, 95/11880, 95/14017, 95/15311, 95/16679, 95/17382, 95/18124, 95/18129, 95/19344, 95/20575, 95/21819, 95/22525, 95/23798, 95/26338, 95/28418, 95/30674, 95/30687, 95/33744, 96/05181, 96/05193, 96/05203, 96/06094, 96/07649, 96/10562, 96/16939, 96/18643, 96/20197, 96/21661, 96/29304, 96/29317, 96/29326, 96/29328, 96/31214, 96/32385, 96/37489, 97/01553, 97/01554, 97/03066, 97/08144, 97/14671, 97/17362, 97/18206, 97/19084, 97/19942 and 97/21702; and in British Patent Publication Nos. 2 266 529, 2 268 931, 2 269 170, 2 269 590, 2 271 774, 2 292 144, 2 293 168, 2 293 169, and 2 302 689. The preparation of such compounds is fully described in the aforementioned patents and publications, which are incorporated herein by reference.

In an embodiment, the neurokinin-1 receptor antagonist for use in conjunction with the compounds of the present invention is selected from: 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)morpholine, or a pharmaceutically acceptable salt thereof, which is described in U.S. Patent No. 5,719,147.

A compound of the instant invention may also be administered with an agent useful in the treatment of anemia. Such an anemia treatment agent is, for example, a continuous erythropoiesis receptor activator (such as epoetin alfa).

A compound of the instant invention may also be administered with an agent useful in the treatment of neutropenia. Such a neutropenia treatment agent is, for example, a hematopoietic growth factor which regulates the production and function of neutrophils such as a human granulocyte colony stimulating factor, (G-CSF). Examples of a G-CSF include filgrastim.

A compound of the instant invention may also be administered with an immunologic-enhancing drug, such as levamisole, isoprinosine and Zadaxin.

Thus, the scope of the instant invention encompasses the use of the instantly claimed compounds in combination with a second compound selected from:

- 5           1)       an estrogen receptor modulator,
- 2)       an androgen receptor modulator,
- 3)       retinoid receptor modulator,
- 4)       a cytotoxic/cytostatic agent,
- 5)       an antiproliferative agent,
- 10       6)       a prenyl-protein transferase inhibitor,
- 7)       an HMG-CoA reductase inhibitor,
- 8)       an HIV protease inhibitor,
- 9)       a reverse transcriptase inhibitor,
- 10)      an angiogenesis inhibitor,
- 15       11)      a PPAR- $\gamma$  agonists,
- 12)      a PPAR- $\delta$  agonists,
- 13)      an inhibitor of inherent multidrug resistance,
- 14)      an anti-emetic agent,
- 15)      an agent useful in the treatment of anemia,
- 20       16)      an agent useful in the treatment of neutropenia,
- 17)      an immunologic-enhancing drug,
- 18)      an inhibitor of cell proliferation and survival signaling,
- 19)      an agent that interferes with a cell cycle checkpoint, and
- 20)      an apoptosis inducing agent.

25           The term "administration" and variants thereof (e.g., "administering" a compound) in reference to a compound of the invention means introducing the compound or a prodrug of the compound into the system of the animal in need of treatment. When a compound of the invention or prodrug thereof is provided in combination with one or more other active agents  
30 (e.g., a cytotoxic agent, etc.), "administration" and its variants are each understood to include concurrent and sequential introduction of the compound or prodrug thereof and other agents.

          As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which  
35 results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The term "therapeutically effective amount" as used herein means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

5           The term "treating cancer" or "treatment of cancer" refers to administration to a mammal afflicted with a cancerous condition and refers to an effect that alleviates the cancerous condition by killing the cancerous cells, but also to an effect that results in the inhibition of growth and/or metastasis of the cancer.

10           In an embodiment, the angiogenesis inhibitor to be used as the second compound is selected from a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP (matrix metalloprotease) inhibitor, an integrin blocker, interferon- $\alpha$ , interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, or an  
15           antibody to VEGF. In an embodiment, the estrogen receptor modulator is tamoxifen or raloxifene.

Also included in the scope of the claims is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with radiation therapy and/or in combination with a compound selected from:

- 20           1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 25           6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor,
- 10) an angiogenesis inhibitor,
- 30           11) PPAR- $\gamma$  agonists,
- 12) PPAR- $\delta$  agonists,
- 13) an inhibitor of inherent multidrug resistance,
- 14) an anti-emetic agent,
- 15) an agent useful in the treatment of anemia,
- 35           16) an agent useful in the treatment of neutropenia,

- 17) an immunologic-enhancing drug,
- 18) an inhibitor of cell proliferation and survival signaling,
- 19) an agent that interferes with a cell cycle checkpoint, and
- 20) an apoptosis inducing agent.

5

And yet another embodiment of the invention is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with paclitaxel or trastuzumab.

10 The invention further encompasses a method of treating or preventing cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with a COX-2 inhibitor.

The instant invention also includes a pharmaceutical composition useful for treating or preventing cancer that comprises a therapeutically effective amount of a compound of Formula I and a compound selected from:

- 15 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 20 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor,
- 10) an angiogenesis inhibitor, and
- 25 11) a PPAR- $\gamma$  agonist,
- 12) a PPAR- $\delta$  agonists;
- 13) an inhibitor of cell proliferation and survival signaling,
- 14) an agent that interferes with a cell cycle checkpoint, and
- 15) an apoptosis inducing agent.

30

The invention further comprises the use of the instant compounds in a method to screen for other compounds that bind to KSP. To employ the compounds of the invention in a method of screening for compounds that bind to KSP kinesin, the KSP is bound to a support, and a compound of the invention (which is a mitotic agent) is added to the assay. Alternatively, the  
35 compound of the invention is bound to the support and KSP is added. Classes of compounds

among which novel binding agents may be sought include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled *in vitro* protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

The determination of the binding of the mitotic agent to KSP may be done in a number of ways. In a preferred embodiment, the mitotic agent (the compound of the invention) is labeled, for example, with a fluorescent or radioactive moiety and binding determined directly. For example, this may be done by attaching all or a portion of KSP to a solid support, adding a labeled mitotic agent (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the amount of the label is that present on the solid support. Various blocking and washing steps may be utilized as is known in the art.

By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g., radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific-binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

In some embodiments, only one of the components is labeled. For example, the kinesin proteins may be labeled at tyrosine positions using  $^{125}\text{I}$ , or with fluorophores.

Alternatively, more than one component may be labeled with different labels; using  $^{125}\text{I}$  for the proteins, for example, and a fluorophor for the mitotic agents.

The compounds of the invention may also be used as competitors to screen for additional drug candidates. "Candidate bioactive agent" or "drug candidate" or grammatical equivalents as used herein describe any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity. They may be capable of directly or indirectly altering the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular proliferation protein binding and/or activity is screened. Screens of this sort may be performed either in the presence or absence of microtubules. In the case where protein binding or activity is screened, preferred embodiments exclude molecules already

known to bind to that particular protein, for example, polymer structures such as microtubules, and energy sources such as ATP. Preferred embodiments of assays herein include candidate agents which do not bind the cellular proliferation protein in its endogenous native state termed herein as "exogenous" agents. In another preferred embodiment, exogenous agents further  
5 exclude antibodies to KSP.

Candidate agents can encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding and lipophilic  
10 binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines,  
15 derivatives, structural analogs or combinations thereof. Particularly preferred are peptides.

Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including  
20 expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification to produce structural analogs.

Competitive screening assays may be done by combining KSP and a drug  
25 candidate in a first sample. A second sample comprises a mitotic agent, KSP and a drug candidate. This may be performed in either the presence or absence of microtubules. The binding of the drug candidate is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of an agent capable of binding to KSP and  
30 potentially modulating its activity. That is, if the binding of the drug candidate is different in the second sample relative to the first sample, the drug candidate is capable of binding to KSP.

In a preferred embodiment, the binding of the candidate agent is determined through the use of competitive binding assays. In this embodiment, the competitor is a binding moiety known to bind to KSP, such as an antibody, peptide, binding partner, ligand, etc. Under

certain circumstances, there may be competitive binding as between the candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

In one embodiment, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to KSP for a time sufficient to allow binding, if present.

5 Incubations may be performed at any temperature which facilitates optimal activity, typically between about 4 and about 40°C.

Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and  
10 the presence or absence of the labeled component is followed, to indicate binding.

In a preferred embodiment, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent is binding to KSP and thus is capable of binding to, and potentially modulating, the activity of KSP. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled,  
15 the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

In an alternative embodiment, the candidate agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate the candidate agent is bound to KSP with a higher affinity. Thus, if the candidate agent is  
20 labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate the candidate agent is capable of binding to KSP.

It may be of value to identify the binding site of KSP. This can be done in a variety of ways. In one embodiment, once KSP has been identified as binding to the mitotic agent, KSP is fragmented or modified and the assays repeated to identify the necessary  
25 components for binding.

Modulation is tested by screening for candidate agents capable of modulating the activity of KSP comprising the steps of combining a candidate agent with KSP, as above, and determining an alteration in the biological activity of KSP. Thus, in this embodiment, the candidate agent should both bind to KSP (although this may not be necessary), and alter its  
30 biological or biochemical activity as defined herein. The methods include both in vitro screening methods and in vivo screening of cells for alterations in cell cycle distribution, cell viability, or for the presence, morphology, activity, distribution, or amount of mitotic spindles, as are generally outlined above.

Alternatively, differential screening may be used to identify drug candidates that  
35 bind to the native KSP, but cannot bind to modified KSP.

Positive controls and negative controls may be used in the assays. Preferably all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the binding of the agent to the protein. Following incubation, all samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

A variety of other reagents may be included in the screening assays. These include reagents like salts, neutral proteins, e.g., albumin, detergents, etc which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

These and other aspects of the invention will be apparent from the teachings contained herein.

### ASSAYS

The compounds of the instant invention described in the Examples were tested by the assays described below and were found to have kinase inhibitory activity. Other assays are known in the literature and could be readily performed by those of skill in the art.

#### I. Kinesin ATPase *In Vitro* Assay

##### **Cloning and expression of human poly-histidine tagged KSP motor domain (KSP(367H))**

Plasmids for the expression of the human KSP motor domain construct were cloned by PCR using a pBluescript full length human KSP construct (Blangy et al., Cell, vol.83, pp1159-1169, 1995) as a template. The N-terminal primer 5'-GCAACGATTAATATGGCGTCGCAGCCAAATTCGTCTGCGAAG (SEQ.ID.NO.: 1) and the C-terminal primer 5'-GCAACGCTCGAGTCAGTGATGATGGTGGTGATGCTGATTCACTTCAGGCTTATTCAATAT (SEQ.ID.NO.: 2) were used to amplify the motor domain and the neck linker region. The PCR products were digested with AseI and XhoI, ligated into the NdeI/XhoI digestion product of pRSETa (Invitrogen) and transformed into E. coli BL21 (DE3).

Cells were grown at 37°C to an OD<sub>600</sub> of 0.5. After cooling the culture to room temperature expression of KSP was induced with 100μM IPTG and incubation was continued



overnight. Cells were pelleted by centrifugation and washed once with ice-cold PBS. Pellets were flash-frozen and stored  $-80^{\circ}\text{C}$ .

### Protein Purification

5 Cell pellets were thawed on ice and resuspended in lysis buffer (50mM K-HEPES, pH 8.0, 250mM KCl, 0.1% Tween, 10mM imidazole, 0.5mM Mg-ATP, 1mM PMSF, 2mM benzimidazole, 1x complete protease inhibitor cocktail (Roche)). Cell suspensions were incubated with 1mg/ml lysozyme and 5mM  $\beta$ -mercaptoethanol on ice for 10 minutes, followed by sonication (3x 30sec). All subsequent procedures were performed at  $4^{\circ}\text{C}$ . Lysates were  
10 centrifuged at 40,000x g for 40 minutes. Supernatants were diluted and loaded onto an SP Sepharose column (Pharmacia, 5ml cartridge) in buffer A (50mM K-HEPES, pH 6.8, 1mM  $\text{MgCl}_2$ , 1mM EGTA, 10 $\mu\text{M}$  Mg-ATP, 1mM DTT) and eluted with a 0 to 750mM KCl gradient in buffer A. Fractions containing KSP were pooled and incubated with Ni-NTA resin (Qiagen) for one hour. The resin was washed three times with buffer B (Lysis buffer minus PMSF and  
15 protease inhibitor cocktail), followed by three 15-minute incubations and washes with buffer B. Finally, the resin was incubated and washed for 15 minutes three times with buffer C (same as buffer B except for pH 6.0) and poured into a column. KSP was eluted with elution buffer (identical to buffer B except for 150mM KCl and 250mM imidazole). KSP-containing fractions were pooled, made 10% in sucrose, and stored at  $-80^{\circ}\text{C}$ .

20 Microtubules are prepared from tubulin isolated from bovine brain. Purified tubulin (> 97% MAP-free) at 1 mg/ml is polymerized at  $37^{\circ}\text{C}$  in the presence of 10  $\mu\text{M}$  paclitaxel, 1 mM DTT, 1 mM GTP in BRB80 buffer (80 mM K-PIPES, 1 mM EGTA, 1 mM  $\text{MgCl}_2$  at pH 6.8). The resulting microtubules are separated from non-polymerized tubulin by ultracentrifugation and removal of the supernatant. The pellet, containing the microtubules, is  
25 gently resuspended in 10  $\mu\text{M}$  paclitaxel, 1 mM DTT, 50  $\mu\text{g/ml}$  ampicillin, and 5  $\mu\text{g/ml}$  chloramphenicol in BRB80.

The kinesin motor domain is incubated with microtubules, 1 mM ATP (1:1  $\text{MgCl}_2$ : Na-ATP), and compound at  $23^{\circ}\text{C}$  in buffer containing 80 mM K-HEPES (pH 7.0), 1 mM EGTA, 1 mM DTT, 1 mM  $\text{MgCl}_2$ , and 50 mM KCl. The reaction is terminated by a 2-10 fold  
30 dilution with a final buffer composition of 80 mM HEPES and 50 mM EDTA. Free phosphate from the ATP hydrolysis reaction is measured via a quinaldine red/ammonium molybdate assay by adding 150  $\mu\text{l}$  of quench C buffer containing a 2:1 ratio of quench A:quench B. Quench A contains 0.1 mg/ml quinaldine red and 0.14% polyvinyl alcohol; quench B contains 12.3 mM ammonium molybdate tetrahydrate in 1.15 M sulfuric acid. The reaction is incubated for 10  
35 minutes at  $23^{\circ}\text{C}$ , and the absorbance of the phospho-molybdate complex is measured at 540 nm.

The compounds 3-1, 4-2, 5-3, 5-4, 7-1, 7-2, 7-3, 8-6a/8-6b, 9-1, 10-2, 11-1, 12-1, 12-2 and 12-3 in the Examples were tested in the above assay and found to have an  $IC_{50} \leq 50\mu M$ .

## II. Cell Proliferation Assay

5 Cells are plated in 96-well tissue culture dishes at densities that allow for logarithmic growth over the course of 24, 48, and 72 hours and allowed to adhere overnight. The following day, compounds are added in a 10-point, one-half log titration to all plates. Each titration series is performed in triplicate, and a constant DMSO concentration of 0.1% is maintained throughout the assay. Controls of 0.1% DMSO alone are also included. Each  
10 compound dilution series is made in media without serum. The final concentration of serum in the assay is 5% in a 200  $\mu L$  volume of media. Twenty microliters of Alamar blue staining reagent is added to each sample and control well on the titration plate at 24, 48, or 72 hours following the addition of drug and returned to incubation at 37°C. Alamar blue fluorescence is analyzed 6-12 hours later on a CytoFluor II plate reader using 530-560 nanometer wavelength  
15 excitation, 590 nanometer emission.

A cytotoxic  $EC_{50}$  is derived by plotting compound concentration on the x-axis and average percent inhibition of cell growth for each titration point on the y-axis. Growth of cells in control wells that have been treated with vehicle alone is defined as 100% growth for the assay, and the growth of cells treated with compounds is compared to this value. Proprietary in-house  
20 software is used to calculate percent cytotoxicity values and inflection points using logistic 4-parameter curve fitting. Percent cytotoxicity is defined as:

$$\% \text{ cytotoxicity} = (\text{Fluorescence}_{\text{control}} - (\text{Flourescence}_{\text{sample}}) \times 100 \times (\text{Fluorescence}_{\text{control}})^{-1}$$

25 The inflection point is reported as the cytotoxic  $EC_{50}$ .

## III. Evaluation of mitotic arrest and apoptosis by FACS

FACS analysis is used to evaluate the ability of a compound to arrest cells in mitosis and to induce apoptosis by measuring DNA content in a treated population of cells. Cells  
30 are seeded at a density of  $1.4 \times 10^6$  cells per  $6\text{cm}^2$  tissue culture dish and allowed to adhere overnight. Cells are then treated with vehicle (0.1% DMSO) or a titration series of compound for 8-16 hours. Following treatment, cells are harvested by trypsinization at the indicated times and pelleted by centrifugation. Cell pellets are rinsed in PBS and fixed in 70% ethanol and stored at 4°C overnight or longer.

For FACS analysis, at least 500,000 fixed cells are pelleted and the 70% ethanol is removed by aspiration. Cells are then incubated for 30 min at 4°C with RNase A (50 Kunitz units/ml) and propidium iodide (50 µg/ml), and analyzed using a Becton Dickinson FACSCaliber. Data (from 10,000 cells) is analyzed using the Modfit cell cycle analysis modeling software (Verity Inc.).

An EC<sub>50</sub> for mitotic arrest is derived by plotting compound concentration on the x-axis and percentage of cells in the G2/M phase of the cell cycle for each titration point (as measured by propidium iodide fluorescence) on the y-axis. Data analysis is performed using the SigmaPlot program to calculate an inflection point using logistic 4-parameter curve fitting. The inflection point is reported as the EC<sub>50</sub> for mitotic arrest. A similar method is used to determine the compound EC<sub>50</sub> for apoptosis. Here, the percentage of apoptotic cells at each titration point (as determined by propidium iodide fluorescence) is plotted on the y-axis, and a similar analysis is carried out as described above.


## VI. Immunofluorescence Microscopy to Detect Monopolar Spindles

Methods for immunofluorescence staining of DNA, tubulin, and pericentrin are essentially as described in Kapoor *et al.* (2000) J. Cell Biol. 150: 975-988. For cell culture studies, cells are plated on tissue culture treated glass chamber slides and allowed to adhere overnight. Cells are then incubated with the compound of interest for 4 to 16 hours. After incubation is complete, media and drug are aspirated and the chamber and gasket are removed from the glass slide. Cells are then permeabilized, fixed, washed, and blocked for nonspecific antibody binding according to the referenced protocol. Paraffin-embedded tumor sections are deparaffinized with xylene and rehydrated through an ethanol series prior to blocking. Slides are incubated in primary antibodies (mouse monoclonal anti-α-tubulin antibody, clone DM1A from Sigma diluted 1:500; rabbit polyclonal anti-pericentrin antibody from Covance, diluted 1:2000) overnight at 4°C. After washing, slides are incubated with conjugated secondary antibodies (FITC-conjugated donkey anti-mouse IgG for tubulin; Texas red-conjugated donkey anti-rabbit IgG for pericentrin) diluted to 15µg/ml for one hour at room temperature. Slides are then washed and counterstained with Hoechst 33342 to visualize DNA. Immunostained samples are imaged with a 100x oil immersion objective on a Nikon epifluorescence microscope using Metamorph deconvolution and imaging software.


## EXAMPLES

Examples provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be illustrative of the invention and not limiting of the reasonable scope thereof.

5

SCHEME 1

## SCHEME 1 (continued)

Step 1: 4-Allyl-4-phenyl-1,3-oxazolidin-2-one (1-4)

- 5 To a suspension of 15.8g (416mmol) of LAH powder in 600 mL of diethyl ether was added 18.3g (90 mmol) of  $\alpha$ -allyl- $\alpha$ -phenylglycine ethyl ester (**1-3**) (prepared according to: Van Betsbrugge et. al. *Tetrahedron*, **1997**, *53*, 9233-9240) in 75 mL of diethyl ether at such a rate as to maintain gentle reflux. After stirring overnight at room temperature, the reaction was carefully quenched with 27 mL of water, followed by 27 mL of 15% NaOH and finally 82 mL of
- 10 water. A quantity of Na<sub>2</sub>SO<sub>4</sub> was added, and the mixture was stirred for 1h. The solids were then filtered off and the solution concentrated. The residue was dissolved in 300 mL of CH<sub>2</sub>Cl<sub>2</sub>, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to provide the amino alcohol as a colorless oil. The amino alcohol (4.5g, 25 mmol) was dissolved in 50 mL of CH<sub>2</sub>Cl<sub>2</sub> and cooled to 0°C. Following the

addition of 5.4 mL (53 mmol) of triethylamine and 4.5g (28 mmol) of 1,1'-carbonyldiimidazole, the mixture was warmed to room temperature and allowed to stir for 4h. The reaction was then dumped into a separatory funnel, washed twice with 1M HCl, water, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to obtain oxazolidinone 1-4 as a colorless oil. Data for 1-4: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.4 – 7.2 (m, 5H), 6.6 (s, 1H), 5.6 – 5.5 (m, 1H), 5.2 (m, 2H), 4.5 (d, 1H), 4.35 (d, 1H), 2.8 (m, 1H), 2.6 (m, 1H) ppm.

**Step 2: Diester (1-5)**

A solution of 68g (334.6 mmol) of 1-4 in 500 mL of CH<sub>2</sub>Cl<sub>2</sub> was cooled to -78°C and ozone was bubbled through the solution until a pale blue color persisted. O<sub>2</sub> was then bubbled through the solution for 15 minutes, followed by 30 minutes with N<sub>2</sub>. At that time, 491 mL (6.7 moles) of dimethyl sulfide was added, and the solution was stirred overnight while slowly coming to room temperature. The volatiles were removed by rotary evaporation to provide a brown oil. This material was suspended in 1L of tBuOH, and 200 mL (1.9 moles) of 2-methyl-2-butene was added. To this solution was then added dropwise a mixture of 160g (1.33 moles) of NaH<sub>2</sub>PO<sub>4</sub> and 70g (774 mmol) of NaClO<sub>2</sub> in 800 mL of H<sub>2</sub>O. After the addition was complete, the mixture was stirred for an additional 4h. After separating the layers, the organic was concentrated by rotary evaporation, the residue was dissolved in EtOAc and placed in a separatory funnel with the aqueous phase from the reaction. After separation, the aqueous phase was extracted 3 x with EtOAc, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to provide ~ 90g of a yellow gum. This residue was suspended in 500 mL of MeOH, and HCl gas was bubbled through the solution until it was nearly refluxing. The flask was then capped and allowed to stir overnight while cooling to room temperature. The volatiles were removed by rotary evaporation, the residue was loaded onto a silica gel column in CH<sub>2</sub>Cl<sub>2</sub>, and eluted with EtOAc/hexanes to provide the methyl ester as a pale orange gum. This residue was dissolved in 500 mL of THF, cooled to 0°C, and 32.6 mL (220.5 mmol) of tert-butyl bromoacetate was added, followed by 10.6g of NaH (264.6 mmol of a 60% suspension). After the mixture was allowed to warm to room temperature and stir overnight, it was quenched with a saturated NH<sub>4</sub>Cl solution, and extracted twice with EtOAc. The combined organic layers were then washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, concentrated, and the residue purified by silica gel chromatography with EtOAc/hexanes to provide 1-5 as a thick pale yellow gum. Data for 1-5: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.4 – 7.3 (m, 5H), 4.65 (d, 1H), 4.55 (d, 1H), 3.9 (d, 1H), 3.65 (s, 3H), 3.5 (d, 1H), 3.35 (d, 1H), 3.2 (d, 1H), 1.4 (s, 9H) ppm. HRMS (ES) calc'd M + Na for C<sub>18</sub>H<sub>23</sub>NO<sub>6</sub>: 372.1423. Found: 372.1412.

**Step 3:** 7a-Phenyldihydro-1H-pyrrolo[1,2-c][1,3]oxazole-3,6(5H)-dione (1-6)

To a solution of 18.6g (53 mmol) of 1-5 in 150 mL of THF at -78°C was added dropwise 58.6 mL (58.6 mmol) of a 1M solution of LiHMDS in THF. After stirring for 1h at that temperature, the cooling bath was removed and the reaction was allowed to warm to room temperature and stir overnight. The mixture was quenched with a saturated NH<sub>4</sub>Cl solution, extracted twice with EtOAc, washed twice with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was dissolved in 60 mL of formic acid and heated at 100°C for 24h. The volatiles were removed under vacuum and the residue was triturated with CH<sub>2</sub>Cl<sub>2</sub>/hexanes/Et<sub>2</sub>O to provide 1-6 as a beige solid. Data for 1-6: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.5 – 7.3 (m, 5H), 4.7 (d, 1H), 4.3 (d, 1H), 4.2 (d, 1H), 3.5 (d, 1H), 3.1 (d, 1H), 2.95 (d, 1H), 2.9 (d, 1H) ppm.

**Step 4:** 6-(2,5-Difluorophenyl)-7a-phenyl-5,7a-dihydro-1H-pyrrolo[1,2-c][1,3]oxazol-3-one (1-7)

To a suspension of 2.2g (10 mmol) of 1-7 in 150 mL of THF at -78°C was added dropwise 12.2 mL (12.2 mmol) of a 1M solution of NaHMDS in THF. After stirring for 30 min, the solution was allowed to warm to 0°C and held there for 1h. The solution was then cooled back down to -78°C and a solution of 4.35g (12.2 mmol) of N-phenylbis(trifluoromethanesulphonimide) in 10 mL of THF was added. The cooling bath was removed and the mixture was allowed to warm to room temperature and stir overnight. The mixture was quenched with a saturated NH<sub>4</sub>Cl solution, extracted twice with EtOAc, washed twice with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was dissolved in 75 mL of DME and 18 mL of water. To this mixture was added 1.29g (30 mmol) of LiCl, 3.2g (30 mmol) of Na<sub>2</sub>CO<sub>3</sub>, and 4.8g (30 mmol) of 2,5-difluorophenylboronic acid. The solution was then degassed with N<sub>2</sub> for 1 minute, followed by the addition of 630 mg (0.5 mmol) of tetrakis(triphenylphosphine) palladium (0). The reaction was heated at 90°C for 3h, cooled to room temperature, diluted with saturated NaHCO<sub>3</sub>, and extracted twice with EtOAc. The combined organic layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, concentrated, and the residue purified by silica gel chromatography with CH<sub>2</sub>Cl<sub>2</sub>/hexanes to provide 1-7 as a white solid. Data for 1-7: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.5 – 7.3 (m, 5H), 7.1 – 6.9 (m, 3H), 6.8 (s, 1H), 4.9 (d, 1H), 4.75 (d, 1H), 4.5 (d, 1H), 4.25 (d, 1H) ppm. HRMS (ES) calc'd M + H for C<sub>18</sub>H<sub>13</sub>F<sub>2</sub>NO<sub>2</sub>: 314.0987. Found: 314.0993.

**Step 5:** 2-({tert-Butyl(dimethyl)silyl}oxy)methyl)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (1-8)

A suspension of 1.75g (5.6 mmol) 1-7 in 15 mL of EtOH and 10 mL of 3 M NaOH was heated at 60°C for 3h, cooled to room temperature and dumped into a separatory funnel with EtOAc and brine. The layers were separated, the aqueous phase was extracted twice with EtOAc, the combined organic phases were washed twice with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to provide a white solid. To this flask was added 30 mL of CH<sub>2</sub>Cl<sub>2</sub>, 1.5g (22.3 mmol) of imidazole and 1.76g (11.7 mmol) of TBSCl, and the resultant suspension was stirred overnight. The reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed twice with water, dried over Na<sub>2</sub>SO<sub>4</sub>, concentrated, and the residue purified by silica gel chromatography with EtOAc/hexanes to provide 1-8 as a white solid. Data for 1-8: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.6 – 7.3 (m, 5H), 7.1 – 6.9 (m, 3H), 6.75 (s, 1H), 4.25 (d, 1H), 4.1 (d, 1H), 3.95 (d, 1H), 3.75 (d, 1H), 0.9 (s, 9H), 0.1 (s, 3H), 0.05 (s, 3H) ppm.

Step 6: Enantiomeric resolution of Intermediate 1-8


Resolution of the enantiomers was carried out chromatographically using a Chiralpak AD<sup>®</sup> 10 x 50cm column with 1% isopropanol in hexanes (with 0.1% diethylamine) at 150 mL/min. Analytical HPLC analysis of the eluent on a 4 x 250mm Chiralpak AD<sup>®</sup> column with 1% isopropanol in hexanes (with 0.1% diethylamine) at 1 mL/min indicated that first eluting, active enantiomer has R<sub>t</sub> = 5.5 min and the second enantiomer has R<sub>t</sub> = 6.9 min.

Step 7: (2S)-2-([*tert*-Butyl(dimethyl)silyl]oxy)methyl-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carbonyl chloride 1-9

To a solution of 1.95g (6.6 mmol) of triphosgene in 25mL of THF at 0°C was added a solution of 1.31g (3.3 mmol) of the first eluting enantiomer of 1-8 and 915 µL (6.6 mmol) of triethylamine in 10 mL of THF. The ice bath was removed and the reaction was allowed to warm to room temperature and stir for 3h. The reaction was then partitioned between water and EtOAc, the organic solution was dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to provide 1-9 as a brown oil. Data for 1-9: HRMS (ES) calc'd M + H for C<sub>24</sub>H<sub>28</sub>ClF<sub>2</sub>NO<sub>2</sub>Si: 464.1619. Found: 464.1625.




## SCHEME 1A

Alternate synthesis of Diester 1-5

To a biphasic mixture of 14.8g (73 mmol) of 1-4 and 110 mL of  $\text{CH}_2\text{Cl}_2$ , 110 mL of  $\text{CH}_3\text{CN}$ , and 320 mL of water was added approximately 200mg of ruthenium(III) chloride hydrate. Sodium periodate (85.6g, 400 mmol) was then added portion-wise over 1h with rapid stirring. After the addition was complete, the reaction was allowed to stir for 4h more at room temperature. The mixture was diluted with 500mL of water and 1.5 L of EtOAc, and the solids were removed by filtration. The filtrate was placed in a separatory funnel, the phases separated, the aqueous phase extracted twice with EtOAc, the combined organic phases washed twice with brine, and dried over  $\text{Na}_2\text{SO}_4$ . Following concentration, the dark brown solid was dissolved in 250 mL of MeOH and  $\text{HCl(g)}$  was slowly passed through the solution at a rate so as not to increase the temperature of the solution above  $35^\circ\text{C}$ . After 5 min, the reaction was capped and allowed to stir at room temperature overnight. The volatiles were then removed on a rotary evaporator, and the residue was purified by silica gel chromatography with EtOAc/hexanes to provide 13.6g (58 mmol) of the methyl ester as a viscous oil. This residue was then dissolved in 200 mL of THF, cooled to  $0^\circ\text{C}$ , and 10.3 mL (70 mmol) of tert-butyl bromoacetate was added, followed by 2.8g of NaH (70 mmol of a 60% suspension). After the mixture was allowed to warm to room temperature and stir overnight, it was quenched with a saturated  $\text{NH}_4\text{Cl}$  solution, and extracted twice with EtOAc. The combined organic layers were then washed with brine,

dried over  $\text{Na}_2\text{SO}_4$ , concentrated, and the residue purified by silica gel chromatography with EtOAc/hexanes to provide 1-5 as a colorless oil.

SCHEME 1B



5

Step 1: Methyl 4-methylene-2-phenylprolinate (1B-2)

An aqueous solution (300 mL) of phenyl glycine methyl ester-HCl (100 g) was neutralized to pH 8 with 10N NaOH. The aqueous solution was extracted with EtOAc (3 X 200 mL). The combined organic extracts were dried over  $\text{MgSO}_4$ , filtered, and concentrated. The residue (56.7 g, 344 mmol) was dissolved in trimethylorthoformate (100 mL) and treated with benzaldehyde (34.9 mL, 36.4 g, 344 mmol). After stirring for 2 h, the reaction was diluted with  $\text{Et}_2\text{O}$  (200 mL) and washed with water (3 X 50 mL). The organic solution was dried over


10

MgSO<sub>4</sub>, filtered, and concentrated. A portion of the imine residue (26.8 g, 100 mmol) was dissolved in dichloromethane (240 mL) and treated with 160 mL of 10N NaOH, methallyl dichloride (50.0 g, 400 mmol), and Bu<sub>4</sub>NHSO<sub>4</sub> (3.59 g). After stirring for 10 h at rt, the reaction was diluted with dichloromethane and the organic solution separated, dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was redissolved in Et<sub>2</sub>O/1N HCl (200 mL/200 mL) and stirred for 2h. The aqueous phase was separated and neutralized with 10N NaOH (to pH 8). The aqueous mixture was extracted with EtOAc (3 x 200 mL). The combined organic solutions were dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was dissolved in water and neutralized (to pH 8). Extraction of this mixture with EtOAc (X 3) followed by drying over MgSO<sub>4</sub>, filtration, and concentration provided crude 1B-2. Purification of this residue by flash chromatography (SiO<sub>2</sub>; 30% EtOAc/hexanes) provided pure 1B-2. Data for 1B-2: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.51 (m, 2H), 7.42 (m, 3H), 5.03 (s, 1H), 4.95 (s, 1H), 3.71 (m, 5H), 3.41 (m, 1H), 2.80 (m, 1H) ppm.

15 Step 2: 7a-Phenyldihydro-1H-pyrrolo[1,2-c][1,3]oxazole-3,6(5H)-dione (1-6)

A suspension of LiAlH<sub>4</sub> (7.14 g, 188 mmol) in THF (500 mL) was cooled to 0°C and treated with a solution of ester 1B-2 (10.2 g, 47 mmol) in THF (50 mL) over 20 min. After stirring for 30 min at 0°C, the reaction was cautiously quenched by the addition of water (7.1 mL), 15% aq NaOH (7.1 mL), and H<sub>2</sub>O (21.3 mL). Solid Na<sub>2</sub>SO<sub>4</sub> was added and the mixture stirred for 40 min. The mixture was filtered and concentrated. The residue (8.2 g, 43.3 mmol) was dissolved in dichloromethane (300 mL) and treated with triethylamine (9.0 mL, 6.5 g, 65.0 mmol) and carbonyldiimidazole (9.14 g, 56.4 mmol). After stirring for 48 h at rt, the reaction was diluted with dichloromethane and washed with 1N HCl and brine. The organic solution was concentrated and not further purified. A solution of the residue 1B-3 (9.2 g, 42.8 mmol) in dichloromethane (200 mL) was cooled to -78°C and ozone was passed through the solution until a blue color persisted. The solution was purged and treated with dimethylsulfide (35 mL). After gradual warming to rt overnight, the solution was concentrated to a yellow solid. Trituration of this solid with Et<sub>2</sub>O provided pure 1-6. Data for 1-6: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.5 – 7.3 (m, 5H), 4.7 (d, 1H), 4.3 (d, 1H), 4.2 (d, 1H), 3.5 (d, 1H), 3.1 (d, 1H), 2.95 (d, 1H), 2.9 (d, 1H) ppm.

## SCHEME 2



**Step 1: Benzyl 3-fluoro-4-oxopiperidine-1-carboxylate (2-2)**

To a solution of 10.0g (43 mmol) of benzyl-4-oxo-1-piperidinecarboxylate in 25 mL of DMF was added 14.3 mL (103 mmol) of triethylamine and then 6.53 mL (52 mmol) of TMSCl. The reaction was heated at 80°C overnight, cooled to room temperature, and then dumped into  
 5 hexanes in a separatory funnel. The mixture was partitioned with saturated aqueous NaHCO<sub>3</sub>, separated, washed with brine, dried over MgSO<sub>4</sub> and concentrated by rotary evaporation. The residue was dissolved in 500mL of CH<sub>3</sub>CN and treated with 16.7g (47 mmol) of Selectfluor. After 90 min the reaction was concentrated to about half the original volume, partitioned between EtOAc and brine, separated, dried over MgSO<sub>4</sub>, filtered, and concentrated by rotary  
 10 evaporation. The residue was loaded onto a silica gel column and eluted with EtOAc/hexanes to provide 2-2 as a colorless oil.

**Step 2: Benzyl 3-fluoro-4-(methylamino)piperidine-1-carboxylate (2-2a)**

15 To a solution of 9.4g (37.5 mmol) of 2-2 in 150 mL of 1,2-dichloroethane was added 37.5 mL (74.9 mmol) of a 2M solution of methylamine in THF and 11.9g (56.2 mmol) of Na(OAc)<sub>3</sub>BH. After stirring for 2h, the reaction was quenched with saturated aqueous K<sub>2</sub>CO<sub>3</sub>, partitioned with EtOAc, separated, and the aqueous phase extracted 3 x EtOAc. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated by rotary  
 20 evaporation. The residue was loaded onto a silica gel column and eluted with 80:10:10 CHCl<sub>3</sub>/EtOAc/MeOH to provide both the *cis* and *trans* isomers of 2-2a as colorless oils. Data for the *trans* isomer of 2-2a, first to elute (confirmed by NOE analysis): <sup>1</sup>HNMR (600 MHz, CD<sub>2</sub>Cl<sub>2</sub>) δ 7.4 – 7.3 (m, 5H), 5.1 (m, 2H), 4.4 – 4.1 (m, 2H), 3.9 (m, 1H), 3.15 - 3.05 (m, 2H), 2.75 (m, 1H), 2.4 (s, 3H), 2.0 (m, 1H), 1.25 (m, 1H) ppm. Data for the *cis* isomer of 2-2a, second  
 25 to elute (confirmed by NOE analysis): <sup>1</sup>HNMR (600 MHz, CD<sub>2</sub>Cl<sub>2</sub>) δ 7.4 – 7.2 (m, 5H), 5.1 (m, 2H), 4.9 – 4.7 (m 1H), 4.4 (m, 1H), 4.15 (m, 1H), 3.1 – 2.9 (m, 2H), 2.6 (m, 1H), 2.4 (s, 3H), 1.8 (m, 1H), 1.6 (m, 1H) ppm. HRMS (ES) calc'd M + H for C<sub>14</sub>H<sub>19</sub>F<sub>1</sub>N<sub>2</sub>O<sub>2</sub>: 267.1504. Found: 267.1500.

**Step 3: Benzyl (3*S*,4*R*)-4-[(*tert*-butoxycarbonyl)(methyl)amino]-3-fluoropiperidine-1-carboxylate (2-3)**


30 To a solution of 7.67g (28.8 mmol) of *cis*-2-2a in 150 mL of CH<sub>2</sub>Cl<sub>2</sub> was added 12.1 mL (86.5 mmol) of triethylamine and 9.44g (43.3 mmol) of di-*tert*-butyl dicarbonate. After stirring for 1h, the reaction was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O, the organic phase was  
 35 washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated by rotary evaporation. The

residue was loaded onto a silica gel column and eluted with EtOAc/hexanes to provide racemic *cis*-2-3 as a white solid. Resolution of the enantiomers was carried out chromatographically using a Chiralpak AD<sup>®</sup> 10 x 50cm column with 20% isopropanol in hexanes (with 0.1% diethylamine) at 150 mL/min. Analytical HPLC analysis of the eluent on a 4 x 250mm Chiralpak AD<sup>®</sup> column with 20% isopropanol in hexanes (with 0.1% diethylamine) at 1 mL/min indicated that first eluting enantiomer (enantiomer of 2-3) has  $R_t$  = 5.90 min and the second enantiomer (2-3) has  $R_t$  = 6.74 min. Data for 2-3: HRMS (ES) calc'd M + Na for C<sub>19</sub>H<sub>27</sub>F<sub>1</sub>N<sub>2</sub>O<sub>4</sub>: 389.1847. Found: 389.1852.

- 10 **Step 4:** *tert*-Butyl [(3*S*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]methylcarbamate (2-4)  
 To a solution of 4.6g (12.6 mmol) of the second eluting enantiomer 2-3 in 150 mL of EtOH was added 29.7 mL (314 mmol) of 1,4-cyclohexadiene and a catalytic amount of 10% Pd on carbon. After stirring overnight, the reaction was filtered through Celite, and concentrated by rotary evaporation. The residue was dissolved in 75 mL of MeOH, 2mL of AcOH and 3.1mL (38 mmol) of 37% aqueous formaldehyde were added, and the mixture was stirred for 1h. At that time, 1.58g (25.1 mmol) of NaCNBH<sub>3</sub> in 10mL of MeOH was added and the reaction was aged for 2h more before being dumped into saturated aqueous NaHCO<sub>3</sub>. After extracting with 3 x CH<sub>2</sub>Cl<sub>2</sub>, the organic phase was washed with water, dried over MgSO<sub>4</sub>, filtered, and concentrated by rotary evaporation to provide 2-4 as a colorless oil. Data for 2-4: HRMS (ES) calc'd M + H for C<sub>12</sub>H<sub>23</sub>FN<sub>2</sub>O<sub>2</sub>: 247.1817. Found: 247.1810.

- Step 5:** (3*S*,4*R*)-3-Fluoro-*N*,1-dimethylpiperidin-4-amine (2-5)  
 To a solution of 3.0g (12.2 mmol) of 2-4 in 100mL of EtOAc was bubbled HCl gas until the solution was warm to the touch. The flask was then capped and stirred for 4h. The volatiles were removed by rotary evaporation, and the residue was triturated with Et<sub>2</sub>O and placed under high vacuum to provide a white solid. This material was mixed with 25 mL of 15% aqueous Na<sub>2</sub>CO<sub>3</sub> and extracted with 5 x 50 mL 2:1 CHCl<sub>3</sub>/EtOH. The organic was concentrated by rotary evaporation with very mild heating, the residue was dissolved in 200 mL of CHCl<sub>3</sub>, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to provide 2-5 as a colorless oil. Data for 2-5: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 4.8 (m, 1H), 3.15 (m, 1H), 2.85 (m, 1H), 2.5 (s, 3H), 2.45 (m, 1H), 2.3 (s, 3H), 2.2 – 2.0 (m, 2H), 1.9 – 1.7 (m, 2H) ppm. HRMS (ES) calc'd M + H for C<sub>7</sub>H<sub>15</sub>FN<sub>2</sub>: 147.1292. Found: 147.1300.

## SCHEME 3




(2S)-4-(2,5-Difluorophenyl)-N-[(3S,4R)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (3-1)

- 5 To a solution of 1.6g (3.45 mmol) of carbamoyl chloride **1-9** in 25mL of THF was added 630mg (4.31 mmol) of amine **2-5**, 1.44 mL (10.3 mmol) of triethylamine, and a catalytic amount of DMAP. After stirring for 24h at room temperature, the reaction was partitioned between EtOAc and saturated aqueous NaHCO<sub>3</sub>, the organic was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated by rotary evaporation. The residue was purified by silica gel
- 10 chromatography with EtOAc/hexanes to provide ~ 1.7g of a pale yellow taffy. This was dissolved in 75 mL of CH<sub>3</sub>CN, 3mL of triethylamine trihydrofluoride was added, and the mixture was stirred for 24h at room temperature. An additional 3 mL of triethylamine trihydrofluoride was added and the reaction was heated at 37°C for an additional 24h. The reaction was then dumped into saturated aqueous NaHCO<sub>3</sub>, extracted 3 x with EtOAc, washed with brine, dried
- 15 over Na<sub>2</sub>SO<sub>4</sub>, and concentrated by rotary evaporation. The residue was purified by silica gel

chromatography with EtOAc - 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O to provide 3-1 as a white solid. Data for 3-1: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.4 – 7.2 (m, 5H), 7.1 – 6.9 (m, 3H), 6.3 (s, 1H), 5.25 (m, 1H), 4.9 (d, 1H), 4.8 (d, 1H), 4.6 (d, 1H), 4.45 (m, 1H), 4.1 – 4.0 (m, 2H), 3.2 – 3.1 (m, 1H), 3.1 (s, 3H), 3.0 (m, 1H), 2.4 – 2.3 (m, 1H), 2.3 (s, 3H), 2.3 – 2.2 (m, 2H), 1.7 (m, 1H) ppm. HRMS (ES) calc'd M + H for C<sub>25</sub>H<sub>28</sub>F<sub>3</sub>N<sub>3</sub>O<sub>2</sub>: 460.2207. Found: 460.2213.

## SCHEME 4




10 (2S)-4-(2,5-Difluorophenyl)-N-[(3R,4S)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (4-2)

This compound was made in a manner identical to that for 3-1, with the exception of incorporation of the first eluting enantiomer of 2-3 from the chiral column described in Scheme 2, Step 3. Data for 4-2: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.4 – 7.2 (m, 5H), 7.1 – 6.9 (m, 3H), 6.3 (s, 1H), 5.4 (bs, 1H), 5.2 (d, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.4 (m, 1H), 4.0 (m, 1H), 3.8 (m, 1H),




3.2 (s, 3H), 3.0 (m, 1H), 2.5 – 2.2 (m, 3H), 2.4 (s, 3H), 1.8 – 1.6 (m, 2H) ppm. HRMS (ES) calc'd M + H for  $C_{25}H_{28}F_3N_3O_2$ : 460.2207. Found: 460.2229.

SCHEME 5



## SCHEME 5 (continued)



(2S)-4-(2,5-Difluorophenyl)-N-[(3R,4R)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (5-3)


- 5 This compound was made in a manner identical to that for 3-1, with the exception that the *trans* isomer of 2-2a was incorporated into the synthesis. Resolution of the enantiomers of “*trans*-2-3” was carried out chromatographically using a Chiralpak AD<sup>®</sup> 10 x 50cm column with 15% EtOH in hexanes (with 0.1% diethylamine) at 150 mL/min. Analytical HPLC analysis of the eluent on a 4 x 250mm Chiralpak AD<sup>®</sup> column with 15% EtOH in hexanes (with 0.1% diethylamine) at 1 mL/min indicated that first eluting enantiomer has  $R_t = 7.30$  min and the second enantiomer has  $R_t = 11.59$  min. The first eluting enantiomer (having unknown absolute stereochemistry) was further processed to provide *trans*-amine 5-1, which was incorporated into the synthesis of 5-3 following a route identical to that explained in Scheme 2. Data for 5-3:
- 15 <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.4 – 7.2 (m, 5H), 7.1 – 6.9 (m, 3H), 6.3 (s, 1H), 5.0 – 4.7 (m, 4H), 4.5 (m, 1H), 4.0 (m, 1H), 3.8 (m, 1H), 3.3 (m, 1H), 2.9 (s, 3H), 2.85 (m, 1H), 2.35 (s, 3H), 2.2 – 1.9 (m, 3H), 1.7 (m, 1H) ppm. HRMS (ES) calc'd M + H for C<sub>25</sub>H<sub>28</sub>F<sub>3</sub>N<sub>3</sub>O<sub>2</sub>: 460.2207. Found: 460.2231. The absolute stereochemistry on the piperidine of compounds 5-3 and 5-4 has not been determined and stereochemistry indicated has been tentatively assigned.

- 20 (2S)-4-(2,5-Difluorophenyl)-N-[(3S,4S)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (5-4)

- This compound was made in a manner identical to that for 5-3, with the exception of incorporation of the second eluting isomer of *trans*-2-3 from the chiral column. Data for 5-4:
- 25 <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.4 – 7.2 (m, 5H), 7.1 – 6.9 (m, 3H), 6.3 (s, 1H), 5.4 (m, 1H), 4.9 – 4.6 (m, 3H), 4.4 (m, 1H), 4.0 (m, 1H), 3.85 (m, 1H), 3.25 (m, 1H), 3.0 (s, 3H), 2.85 (m, 1H), 2.35

(s, 3H), 2.2 – 1.9 (m, 4H) ppm. HRMS (ES) calc'd M + H for  $C_{25}H_{28}F_3N_3O_2$ : 460.2207. Found: 460.2229.

**SCHEME 6**



**Step 1:** (2S,4S)-tert-Butyl 4-hydroxy-2-phenylpyrrolidine-1-carboxylate (6-2)

To a flame dried flask equipped with stir bar was added tert-butyl (2S,4S)-4-  
 {[tert-butyl(dimethyl)silyl]oxy}-2-phenylpyrrolidine-1-carboxylate (6-1, prepared from (S)-(-)-4-  
 chloro-3-hydroxybutyronitrile by the method of Maeda, *et al Synlett* 2001, 1808-1810, 7.8 g,  
 20.7 mmol) and anhydrous acetonitrile (20.0 mL). The resulting solution was treated with  
 triethylamine trihydrofluoride (10.1 mL, 62.0 mmol) while stirring under N<sub>2</sub>. The reaction  
 stirred 12 hours at 40°C. The reaction was then diluted with EtOAc (100 mL) and poured into  
 5% aq. NaHCO<sub>3</sub>. Following cessation of gas evolution, the organic layer was washed three  
 additional times with 5% aq. NaHCO<sub>3</sub>. The organic layer was dried over magnesium sulfate,  
 filtered and concentrated to provide crude product. Recrystallization was effected from  
 EtOAc/hexanes to provide (2S,4S)-tert-butyl 4-hydroxy-2-phenylpyrrolidine-1-carboxylate (6-2)  
 as a white crystalline solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) rotamers δ 7.38-7.18 (m, 5H), 4.90 (m,  
 1H), 4.42 (m, 1H), 3.88 (m, 1H), 3.56 (dd, *J* = 11.5, 4.0 Hz, 1H), 2.60 (m, 1H), 2.03 (m, 1H),  
 1.50 and 1.20 (br s, 9H); MS 208.0 found, 208.1 (M – C(CH<sub>3</sub>)<sub>3</sub>) required.

**Step 2:** (2S)-tert-Butyl 4-oxo-2-phenylpyrrolidine-1-carboxylate (6-3)

To a flame dried flask equipped with stir bar was added 150 mL anhydrous  
 dichloromethane which was cooled to –78°C. Oxalyl chloride (3.8 mL, 44 mmol) and DMSO  
 (4.8 mL, 61 mmol) were added sequentially and the reaction stirred for 10 minutes. (2S,4S)-tert-  
 Butyl 4-hydroxy-2-phenylpyrrolidine-1-carboxylate (6-2, 2.28 g, 8.73 mmol) in 10 mL anhydrous  
 dichloromethane was added dropwise and stirred 1 hour at –78°C. Triethylamine (12 mL,  
 87mmol) was added and the reaction was warmed to 0°C over 1 hour. Upon completion, the  
 reaction was washed with 5% NaHCO<sub>3</sub>, brine and dried over MgSO<sub>4</sub>. The organic layer was  
 concentrated to provide crude (2S)-tert-butyl 4-oxo-2-phenylpyrrolidine-1-carboxylate (6-3).  
 Recrystallization was effected with EtOAc/hexanes. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.35 (m,  
 3H), 7.17 (m, 2H), 5.38 (m, 1H), 4.08 (d, *J* = 19.5 Hz, 1H), 3.90 (d, *J* = 19.3 Hz, 1H), 3.13 (dd, *J*  
 = 18.8, 9.8 Hz, 1H), 2.58 (dd, *J* = 18.6, 2.4 Hz, 1H), 1.40 (br s, 9H); MS 206.0 found, 206.1 (M  
 – C(CH<sub>3</sub>)<sub>3</sub>) required.

**Step 3:** (2S)-tert-Butyl 2-phenyl-4-[[trifluoromethyl)sulfonyl]oxy]-2,5-dihydro-1H-pyrrole-1-carboxylate (6-4)

To a flame dried flask equipped with stir bar was added ketone (2S)-tert-butyl 4-  
 oxo-2-phenylpyrrolidine-1-carboxylate (6-3, 0.16 g, 0.62 mmol) and anhydrous THF (2 mL).  
 The resulting solution was cooled to –78°C, and treated dropwise with lithium  
 hexamethyldisilylamide (LHMDS, 0.68 mL, 1M in THF, 0.68 mmol). The reaction stirred 1

hour at  $-78^{\circ}\text{C}$ , and N-(5-chloropyridin-2-yl)-1,1,1-trifluoro-N-  
 [(trifluoromethyl)sulfonyl]methanesulfonamide (0.27 g, 0.68 mmol) was added neat in one  
 portion. The reaction was allowed to warm to  $0^{\circ}\text{C}$  and stirred 4 hours total. The reaction was  
 diluted with  $\text{Et}_2\text{O}$  (10 mL) and washed successively with  $\text{H}_2\text{O}$  (10 mL) and brine (10 mL). The  
 organic layer was dried over  $\text{MgSO}_4$ , filtered and concentrated. The crude residue was purified  
 by flash column chromatography (0-20% EtOAc/hexanes gradient, 15 min) to provide (2S)-tert-  
 butyl 2-phenyl-4-[[[(trifluoromethyl)sulfonyl]oxy]-2,5-dihydro-1H-pyrrole-1-carboxylate (6-4).  
 $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ) major rotamer:  $\delta$  7.30 (m, 5H), 5.72 (m, 1H), 5.48 (m, 1H), 4.42  
 (m, 2H), 1.18 (s, 9H); MS 379.0 found 379.1 ( $\text{M} - \text{CH}_3$ ) required.

**Step 4:** (2S)-4-(2,5-Difluorophenyl)-2-phenyl-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-  
 carboxamide (6-5)


To a flame dried flask equipped with stir bar was added (2S)-tert-butyl 2-phenyl-  
 4-[[[(trifluoromethyl)sulfonyl]oxy]-2,5-dihydro-1H-pyrrole-1-carboxylate (6-4, 0.250 g, 0.636  
 mmol), 2,5-difluorophenyl boronic acid (0.251 g, 1.59 mmol),  $\text{Na}_2\text{CO}_3$  (0.202 g, 1.91 mmol),  
 and LiCl (0.081 g, 1.91 mmol). The solids were dissolved in 20 mL 4:1 DME/ $\text{H}_2\text{O}$  and degassed  
 with nitrogen.  $\text{Pd}(\text{PPh}_3)_4$  (0.037 g, 0.032 mmol) was added and the reaction was sealed under  
 nitrogen and heated to  $90^{\circ}\text{C}$  for 2 hours. Upon completion, the reaction was partitioned between  
 5% aq.  $\text{NaHCO}_3$  and EtOAc (3 x 50 mL), and the combined organic layers were dried over  
 $\text{MgSO}_4$ . Following filtration, the organic layer was concentrated and purified via flash column  
 chromatography ( $\text{SiO}_2$ , 0-20% EtOAc/hexanes gradient) to provide (2S)-tert-butyl 4-(2,5-  
 difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (6-5).

**Step 5:** 1-[[[(2S)-4-(2,5-Difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-  
 3-methyl-1H-imidazol-3-ium (6-6)

To a flame-dried flask equipped with stir bar under nitrogen was charged 6-5  
 (0.63 g, 1.75 mmol) and anhydrous  $\text{CH}_2\text{Cl}_2$  (10 mL). The resulting solution was treated with  
 trifluoroacetic acid (5 mL) and stirred 1.5 hours at  $25^{\circ}\text{C}$ . Upon completion, the reaction was  
 concentrated, taken up in  $\text{CH}_2\text{Cl}_2$  (50 mL) and washed with 5%  $\text{NaHCO}_3$  (50 mL). The organic  
 layer was dried ( $\text{MgSO}_4$ ), filtered and concentrated under reduced pressure. The resulting free-  
 amine was dissolved in anhydrous THF (10 mL) and treated with carbonyl diimidazole (0.31 g,  
 1.93 mmol). The resulting solution was refluxed for 4 hours until completion. The reaction was  
 concentrated, taken up in EtOAc (50 mL) and washed with  $\text{H}_2\text{O}$  and brine. The combined  
 organic layers were dried ( $\text{MgSO}_4$ ) and concentrated. The crude acyl imidazole was dissolved in  
 anhydrous  $\text{CH}_3\text{CN}$  and treated with MeI (2.2 mL, 36 mmol). The resulting solution was stirred at

25°C overnight. Upon completion, the reaction was concentrated to give **6-6** as an orange colored solid: LRMS  $m/z$  (M+H) 365.9 found, 366.1 required.

SCHEME 7




5

(2S)-4-(2,5-Difluorophenyl)-N-[(3S,4R)-3-fluoro-1-methylpiperidin-4-yl]-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (**7-1**)

To a solution of 75mg (0.15 mmol) of **6-6** in 1mL of THF was added 40mg (0.18 mmol) of the bis-HCl salt of amine **2-5** and 106  $\mu$ L (0.76 mmol) of triethylamine. After stirring for 72h at room temperature, the reaction was partitioned between  $\text{CH}_2\text{Cl}_2$  and saturated aqueous  $\text{NaHCO}_3$ , the organic was washed with brine, dried over  $\text{MgSO}_4$ , and concentrated by rotary evaporation. The residue was purified by silica gel chromatography with 80:10:10


10

CHCl<sub>3</sub>/EtOAc/MeOH to provide 7-1 as a white solid. Data for 7-1: HRMS (ES) calc'd M + H for C<sub>24</sub>H<sub>26</sub>F<sub>3</sub>N<sub>3</sub>O: 430.2101. Found: 430.2116.



5 (2S)-4-(2,5-Difluorophenyl)-N-[(3R,4S)-3-fluoro-1-methylpiperidin-4-yl]-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (7-2)


This compound was made in an analogous way as was 7-1, substituting 4-1 as starting material. Data for 7-2: HRMS (ES) calc'd M + H for C<sub>24</sub>H<sub>26</sub>F<sub>3</sub>N<sub>3</sub>O: 430.2101. Found: 430.2119.









10 (2S)-4-(2,5-Difluorophenyl)-N-[(3R,4R)-3-fluoro-1-methylpiperidin-4-yl]-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (7-3)









This compound was made in an analogous way as was 7-1, substituting 5-1 as starting material. Data for 7-3: HRMS (ES) calc'd M + H for C<sub>24</sub>H<sub>26</sub>F<sub>3</sub>N<sub>3</sub>O: 430.2101. Found: 430.2115.








The following compounds are prepared by simple modifications of the procedures illustrated in Schemes 1-7 and Schemes G-N, but substituting the appropriately substituted reagents for those utilized in the Schemes.



















R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H

















R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
Me	Me	Me	F	H
Me		Me	F	H
Me		Me	F	H


R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H


$R_1$	$R_2$	$R_3$	$R_4$	$R_5$
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me	CH <sub>2</sub> OH	Me	Cl	H
Me	CH <sub>2</sub> OH	Me	Br	H
Me	CH <sub>2</sub> OH	Me	CN	H
Me	CH <sub>2</sub> OH	Me	Me	H
Me	CH <sub>2</sub> OH	Me	CF <sub>3</sub>	H
Me	CH <sub>2</sub> OH	Me	NO <sub>2</sub>	H
Me	CH <sub>2</sub> OH	Me	F	OH
Me	CH <sub>2</sub> OH	Me	F	NH <sub>2</sub>
Me	CH <sub>2</sub> OH	Me	F	F
Me	CH <sub>2</sub> OH	Me	F	SH

SCHEME 8

1.  $\text{SO}_3$ -pyridine, TEA,  
DMSO/ $\text{CH}_2\text{Cl}_2$

2.  $\text{MeNH}_2$ ,  $\text{NaCNBH}_3$ ,  
MeOH/AcOH  
3.  $\text{BOC}_2\text{O}$ , TEA,  $\text{CH}_2\text{Cl}_2$   
4.  $\text{LiBH}_4$ , THF/MeOH





1. Pd/C, 1,4-cyclohexadiene,  
EtOH

2.  $\text{CH}_2\text{O}$ ,  $\text{NaCNBH}_3$ ,  
MeOH/AcOH



## SCHEME 8 (continued)



benzyl 4-[(*tert*-Butoxycarbonyl)(methyl)amino]-2-(hydroxymethyl)piperidine-1-carboxylate (8-2)

- 5 To a solution of 2.0 g (6.8 mmol) of 8-1 (reported in: S. J. Hayes, T. C. Malone, G. Johnson *J. Org. Chem.* 1991, 56, 4084-4086) in 100 mL of  $\text{CH}_2\text{Cl}_2$  was added 3.33 mL (23.9 mmol) of triethylamine, followed by dropwise addition of 2.44g (15.4 mmol) of  $\text{SO}_3$ -pyridine in

50 mL of DMSO. After stirring for 5h at room temperature, the mixture was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O, the phases were separated, washed with 2 x 1M HCl, saturated aqueous NaHCO<sub>3</sub>, brine, dried over MgSO<sub>4</sub>, and concentrated to provide the ketone. To 2.1g (7.2 mmol) of this ketone dissolved in 35 mL of MeOH was added 1mL of AcOH and 14.4 mL (28.9 mmol) of a 2M solution of MeNH<sub>2</sub> in MeOH. After stirring for 1h, 910mg (14.4 mmol) of NaCNBH<sub>3</sub> in 5 mL of MeOH was added and the reaction was stirred overnight. The reaction was quenched with saturated aqueous NH<sub>4</sub>Cl, extracted with CH<sub>2</sub>Cl<sub>2</sub>, washed with NaHCO<sub>3</sub>, H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, and concentrated to provide 2.0g (7.2 mmol) of the amine. This material was dissolved in 25mL of CH<sub>2</sub>Cl<sub>2</sub>, 1.78g (8.2 mmol) of di-*tert*-butyl dicarbonate and 1.8 mL (12.9 mmol) of triethylamine were added, and the resultant mixture was stirred for 72h. The reaction was then partitioned between CH<sub>2</sub>Cl<sub>2</sub> and saturated aqueous NaHCO<sub>3</sub>, separated, washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, and concentrated. The residue was purified by silica gel chromatography with EtOAc/hexanes to provide 1.85g (4.6 mmol) of a mixture of diastereomers. This was dissolved in 20 mL of THF and 1mL of MeOH, cooled to 0°C, and 496mg (22.8 mmol) of LiBH<sub>4</sub> was added. After warming to room temperature and stirring overnight, the reaction was quenched with saturated aqueous NH<sub>4</sub>Cl, extracted with EtOAc, washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, and concentrated to provide 8-2 as a colorless gum. Data for 8-2: LRMS (ES) calc'd M + H for C<sub>20</sub>H<sub>30</sub>N<sub>2</sub>O<sub>5</sub>: 379. Found: 379.

20 *tert*-Butyl [2-(hydroxymethyl)-1-methylpiperidin-4-yl]methylcarbamate (8-3)

To 1.08g (2.9 mmol) of 8-2 in 30 mL of EtOH was added 7mL (74 mmol) of 1,4-cyclohexadiene and a catalytic amount of 10% Pd on carbon. After stirring overnight, the reaction was filtered through Celite, concentrated by rotary evaporation, and dissolved in 25mL of MeOH. To this was added 2mL of AcOH, and 700 µL (8.6 mmol) of 37% aqueous formaldehyde. After stirring overnight, 540mg (8.6 mmol) of NaCNBH<sub>3</sub> in 5 mL of MeOH was added and the reaction was stirred for 1h more. The solvents were removed by rotary evaporation, the residue was partitioned between EtOAc and aqueous NaHCO<sub>3</sub>, the organic phase was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The residue was taken up in CH<sub>2</sub>Cl<sub>2</sub>, filtered and concentrated to provide 8-3 as a colorless oil. Data for 8-3: LRMS (ES) calc'd M + H for C<sub>13</sub>H<sub>26</sub>N<sub>2</sub>O<sub>3</sub>: 259. Found: 259.

*tert*-Butyl [2-(fluoromethyl)-1-methylpiperidin-4-yl]methylcarbamate (8-4) and *tert*-butyl (6-fluoro-1-methylazepan-4-yl)methylcarbamate (8-5)

To a solution of 350 µL (2.6 mmol) of (diethylamino)sulfur trifluoride (DAST) in 15mL of CH<sub>2</sub>Cl<sub>2</sub> at -78°C was added 520mg (2.0 mmol) of 8-3 in 5 mL of CH<sub>2</sub>Cl<sub>2</sub>. The reaction


was allowed to slowly warm to room temperature with stirring overnight, and was then quenched with ice water. The mixture was partitioned with additional  $\text{CH}_2\text{Cl}_2$  and a small amount of 3M KOH, the organic phase was washed with water, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. The residue was loaded onto a silica gel column and eluted with EtOAc - 20:1:1 EtOH/ $\text{NH}_4\text{OH}$ / $\text{H}_2\text{O}$ .








- 5 The first product to elute was the ring enlarged product 8-5 as a colorless oil, and the second to elute was 8-4 as a colorless oil. Both 8-5 and 8-4 were isolated as enantiomeric mixtures of the *trans* diastereomer. The structures were confirmed by extensive 1D and 2D NMR spectroscopy. Data for 8-5:  $^1\text{H}$ NMR (600 MHz,  $\text{CD}_2\text{Cl}_2$ )  $\delta$  4.75 (m, 1H), 4.1 – 3.9 (m, 1H), 2.9 – 2.7 (m, 3H), 2.75 (s, 3H), 2.4 (s, 3H), 2.35 (m, 1H), 2.1 – 1.7 (m, 4H), 1.4 (s, 9H) ppm. Data for 8-4:  $^1\text{H}$ NMR (500 MHz,  $\text{CD}_2\text{Cl}_2$ )  $\delta$  4.8 – 4.5 (m, 2H), 4.1 – 4.0 (m, 1H), 3.2 (m, 1H), 2.75 (m, 2H), 2.7 (s, 3H), 2.45 (s, 3H), 1.9 – 1.5 (m, 4H), 1.45 (s, 9H) ppm.








- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(2*R*,4*R*)-2-(fluoromethyl)-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide (8-6a) and (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(2*S*,4*S*)-2-(fluoromethyl)-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide (8-6b)








- A solution of 80mg (0.31 mmol) of 8-4 in 30 mL of EtOAc was saturated with HCl gas and allowed to stir 1h at room temperature. The reaction was then concentrated by rotary evaporation and the resulting white solid was suspended in 1mL of THF. To this was added 156mg (0.34 mmol) of 1-9, 268  $\mu\text{L}$  (1.54 mmol) of diisopropylethylamine and a catalytic amount of DMAP. After heating at 50  $^\circ\text{C}$  overnight, 500  $\mu\text{L}$  of trifluoroacetic acid was added and stirring was continued an additional 1h at room temperature before being quenched with saturated aqueous  $\text{NaHCO}_3$ . The mixture was partitioned with  $\text{CH}_2\text{Cl}_2$ , the organic phase was washed with brine, dried over  $\text{MgSO}_4$ , and concentrated. The residue was loaded onto a silica gel column and eluted with  $\text{CHCl}_3$  - 80:10:10  $\text{CHCl}_3$ /EtOAc/MeOH to provide 8-6a and 8-6b as an inseparable mixture - a colorless gum. Data for 8-6a/8-6b: HRMS (ES) calc'd M + H for  $\text{C}_{26}\text{H}_{30}\text{F}_3\text{N}_3\text{O}_2$ : 474.2363. Found: 474.2374.











- The following compounds are prepared by simple modifications of the procedures illustrated in Schemes 1-8 and Schemes G-N, but substituting the appropriately substituted reagents for those utilized in the Schemes.









R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H







R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H









R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
Me	Me	Me	F	H
Me		Me	F	H
Me		Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H




R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H

$R_1$	$R_2$	$R_3$	$R_4$	$R_5$
Me	$\text{CH}_2\text{OH}$		F	H
Me	$\text{CH}_2\text{OH}$		F	H
Me	$\text{CH}_2\text{OH}$		F	H
Me	$\text{CH}_2\text{OH}$		F	H
Me	$\text{CH}_2\text{OH}$		F	H
Me	$\text{CH}_2\text{OH}$		F	H
Me	$\text{CH}_2\text{OH}$		F	H
Me	$\text{CH}_2\text{OH}$		F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me	CH <sub>2</sub> OH	Me	Cl	H
Me	CH <sub>2</sub> OH	Me	Br	H
Me	CH <sub>2</sub> OH	Me	CN	H
Me	CH <sub>2</sub> OH	Me	Me	H
Me	CH <sub>2</sub> OH	Me	CF <sub>3</sub>	H
Me	CH <sub>2</sub> OH	Me	NO <sub>2</sub>	H
Me	CH <sub>2</sub> OH	Me	F	OH
Me	CH <sub>2</sub> OH	Me	F	NH <sub>2</sub>
Me	CH <sub>2</sub> OH	Me	F	F
Me	CH <sub>2</sub> OH	Me	F	SH


## SCHEME 9



(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methyl-1-oxidopiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide (**9-1**)

- 5 To a solution of 20mg (0.044 mmol) of **3-1** in 1 mL of CH<sub>2</sub>Cl<sub>2</sub> at 0°C was added 11mg (~ 0.048 mmol) of mCPBA. The ice-bath was removed and the reaction was stirred for 30 min. The mixture was partitioned with EtOAc and saturated aqueous NaHCO<sub>3</sub>, separated, washed with H<sub>2</sub>O, brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated by rotary evaporation. The residue
- 10 was loaded onto a silica gel column and eluted with EtOAc- 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O to provide **9-1** as a white foam. Data for **9-1**: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.4 – 7.2 (m, 5H), 7.1 – 6.9 (m, 3H), 6.25 (s, 1H), 5.0 – 4.9 (m, 2H), 4.7 (m, 1H), 4.5 (m, 1H), 4.3 – 4.2 (m, 1H), 4.0 (m, 2H), 3.7 (m, 1H), 3.4 – 3.3 (m, 2H), 3.3 (s, 3H), 3.2 (s, 3H), 2.5 – 1.9 (bs, 2H), 1.8 (m, 1H) ppm. HRMS (ES) calc'd M + H for C<sub>25</sub>H<sub>28</sub>F<sub>3</sub>N<sub>3</sub>O<sub>3</sub>: 476.2156. Found: 476.2165.

## SCHEME 10



**Step 1:** Benzyl (3*S*,4*R*)-4-[[[(2*S*)-4-(2,5-difluorophenyl)-2-(hydroxymethyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]carbonyl](methylamino)-3-fluoropiperidine-1-carboxylate (**10-1**):


To a solution of 885mg (2.4 mmol) of **2-3** in 12 mL of EtOAc at 0°C was added 12 mL (48 mmol) of a 4M solution of HCl in dioxane. The ice-bath was removed, stirring was continued for 2h, and then the volatiles were removed by rotary evaporation. The residue was partitioned between EtOAc and saturated aqueous NaHCO<sub>3</sub> with 5mL of 1M NaOH, separated, and the aqueous phase extracted 2 x EtOAc. The combined organic extracts were washed again with NaHCO<sub>3</sub>, brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to provide 590mg (2.2 mmol) of the amine. To 215mg (0.81 mmol) of this material in 5 mL of THF was added 340mg (0.73 mmol)

of 1-9, 306  $\mu\text{L}$  (2.2 mmol) of triethylamine, and a catalytic amount of DMAP. After stirring overnight, the reaction was judged only ~ 40% complete by LC/MS, so an additional portion of DMAP was added and stirring was continued overnight. The reaction was then partitioned between EtOAc and saturated aqueous  $\text{NaHCO}_3$ , the organic phase was washed with  $\text{NaHCO}_3$ ,  
 5  $\text{H}_2\text{O}$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. The residue was purified by silica gel chromatography with EtOAc/hexanes to provide 10-1 as a colorless oil. Data for 10-1: LRMS (ES) calc'd M + H for  $\text{C}_{38}\text{H}_{46}\text{F}_3\text{N}_3\text{O}_4\text{Si}$ : 694.4. Found: 694.3.

10 Step 2: (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoropiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide (10-2):

To a solution of 405mg (0.58 mmol) of 10-1 in 4 mL of EtOH was added 1.38 mL (14.6 mmol) of 1,4-cyclohexadiene, 100mg of 10% palladium on carbon, and then stirred  
 15 overnight. The reaction was filtered through Celite, concentrated, and the residue was dissolved in 3 mL of  $\text{CH}_2\text{Cl}_2$ . To this was added 3mL of trifluoroacetic acid, stirring was maintained for 1h, and then the mixture was partitioned between EtOAc and saturated aqueous  $\text{NaHCO}_3$  plus 25mL of 5% aqueous  $\text{Na}_2\text{CO}_3$ , the organic phase was washed with  $\text{NaHCO}_3$  again,  $\text{H}_2\text{O}$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated by rotary evaporation. The residue was loaded onto a silica  
 20 gel column and eluted with EtOAc – 20:1:1 EtOH/ $\text{NH}_4\text{OH}$ / $\text{H}_2\text{O}$  to provide 10-2 as a white solid. There was an impurity of ~ 5% subsequently removed by reverse phase chromatography with 95:5 to 5:95  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  (both with 0.1% TFA). The fractions were basified with  $\text{NaHCO}_3$  to provide pure B-2 as a white solid. Data for 10-2:  $^1\text{H}$ NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.4 – 7.2 (m, 5H), 7.1 – 6.9 (m, 3H), 6.3 (s, 1H), 5.2 (bs, 1H), 4.9 (m, 1H), 4.8 – 4.6 (m, 2H), 4.45 (m, 1H), 4.2  
 25 – 4.1 (m, 1H), 4.0 (m, 1H), 3.3 – 3.2 (m, 2H), 3.1 (s, 3H), 2.85 – 2.7 (m, 2H), 2.1 (m, 1H), 1.7 (m, 2H) ppm. HRMS (ES) calc'd M + H for  $\text{C}_{24}\text{H}_{26}\text{F}_3\text{N}_3\text{O}_2$ : 446.2050. Found: 446.2059.




## SCHEME 11



(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-isopropylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide (**11-1**)

- 5           To a solution of 38mg (0.085 mmol) of **10-2** in 1 mL of 1,2-dichloroethane was added 20  $\mu$ L (0.34 mmol) of acetic acid, 25  $\mu$ L (0.34 mmol) of acetone, and 27 mg (0.13 mmol) of Na(OAc)<sub>3</sub>BH. After stirring overnight, the reaction was partitioned between EtOAc and saturated aqueous NaHCO<sub>3</sub>, the organic phase was washed again with NaHCO<sub>3</sub>, H<sub>2</sub>O, brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated by rotary evaporation. The residue was purified by reverse
- 10 phase chromatography with 95:5 to 5:95 H<sub>2</sub>O/CH<sub>3</sub>CN (both with 0.1% TFA). The fractions were basified with NaHCO<sub>3</sub> to provide **11-1** as a colorless film. Data for **11-1**: <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.4 – 7.2 (m, 5H), 7.1 – 6.9 (m, 3H), 6.3 (s, 1H), 5.3 (m, 1H), 4.95 – 4.8 (m, 2H), 4.6 (m, 1H), 4.45 (m, 1H), 4.1 – 4.0 (m, 2H), 3.2 (m, 1H), 3.15 (s, 3H), 3.0 (m, 1H), 2.8 (m, 1H), 2.45 – 2.25 (m, 3H), 1.75 (m, 1H), 1.1 – 1.0 (m, 6H) ppm. HRMS (ES) calc'd M + H for
- 15 C<sub>27</sub>H<sub>32</sub>F<sub>3</sub>N<sub>3</sub>O<sub>2</sub>: 488.2519. Found: 488.2517.

## SCHEME 12





(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*S*)-3-fluoropiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide (12-1)

---

This compound was made in a manner identical to that for 10-2, with the exception of incorporation of the enantiomer of 2-3. Data for 12-1: HRMS (ES) calc'd M + H for C<sub>24</sub>H<sub>26</sub>F<sub>3</sub>N<sub>3</sub>O<sub>2</sub>: 446.2050. Found: 446.2069.

(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*R*)-3-fluoropiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide (12-2)

---

This compound was made in a manner identical to that for 10-2, with the exception that the *trans* isomer of 2-2a was incorporated into the synthesis. Resolution of the enantiomers of "*trans*-2-3" was carried out chromatographically using a Chiralpak AD<sup>®</sup> 10 x 50cm column with 15% EtOH in hexanes (with 0.1% diethylamine) at 150 mL/min. Analytical HPLC analysis of the eluent on a 4 x 250mm Chiralpak AD<sup>®</sup> column with 15% EtOH in hexanes (with 0.1% diethylamine) at 1 mL/min indicated that first eluting enantiomer has R<sub>t</sub> = 7.30 min and the second enantiomer has R<sub>t</sub> = 11.59 min. The first eluting enantiomer (having unknown absolute stereochemistry) was further processed to provide the *trans* amine, which was incorporated into the synthesis of 12-2 following a route identical to that explained in Scheme 10. Data for 12-2: HRMS (ES) calc'd M + H for C<sub>24</sub>H<sub>26</sub>F<sub>3</sub>N<sub>3</sub>O<sub>2</sub>: 446.2050. Found: 446.2069. The absolute stereochemistry on the piperidine of compounds 12-2 and 12-3 has not been determined and stereochemistry indicated has been tentatively assigned.


(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*S*)-3-fluoropiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide (12-3)

---

This compound was made in a manner identical to that for 10-2, with the exception of incorporation of the second eluting isomer of "*trans*-2-3" from the chiral column. Data for 12-3: HRMS (ES) calc'd M + H for C<sub>24</sub>H<sub>26</sub>F<sub>3</sub>N<sub>3</sub>O<sub>2</sub>: 446.2050. Found: 446.2068.

## WHAT IS CLAIMED IS:

1. A compound of Formula I:



5

or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

- 10 a is 0 or 1;  
 b is 0 or 1;  
 m is 0, 1, or 2;  
 n is 0, 1, 2 or 3;  
 r is 0 or 1;  
 15 s is 0 or 1;  
 t is 0, 1 or 2;  
 u is 0 or 1;

- 20 R<sup>1</sup> and R<sup>2</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>3</sup> is selected from:

- 1) hydrogen;

- 2) C<sub>1</sub>-C<sub>10</sub> alkyl;
  - 3) C<sub>1</sub>-C<sub>10</sub> alkyl-O-R<sup>d</sup>,
  - 4) C<sub>2</sub>-C<sub>10</sub> alkenyl-O-R<sup>d</sup>,
  - 5) C<sub>2</sub>-C<sub>10</sub> alkynyl-O-R<sup>d</sup>,
  - 5 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-O-R<sup>d</sup>,
  - 7) C<sub>1</sub>-C<sub>10</sub> alkyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
  - 8) C<sub>2</sub>-C<sub>10</sub> alkenyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
  - 9) C<sub>2</sub>-C<sub>10</sub> alkynyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
  - 10 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
  - 11) C<sub>1</sub>-C<sub>10</sub> alkyl-S(O)<sub>m</sub>-R<sup>d</sup>,
  - 12) C<sub>2</sub>-C<sub>10</sub> alkenyl-S(O)<sub>m</sub>-R<sup>d</sup>,
  - 13) C<sub>2</sub>-C<sub>10</sub> alkynyl-S(O)<sub>m</sub>-R<sup>d</sup>,
  - 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-S(O)<sub>m</sub>-R<sup>d</sup>,
- 15 said alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more substituents selected from R<sup>6</sup>;

R<sup>4</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 20 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) CO<sub>2</sub>H,
- 4) halo,
- 5) CN,
- 6) OH,
- 25 7) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 8) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>8</sup>R<sup>9</sup>,
- 9) S(O)<sub>m</sub>R<sup>a</sup>,
- 10) S(O)<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, and
- 11) -OPO(OH)<sub>2</sub>;

30 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>5</sup> is selected from:

- 1) hydrogen;
- 2)  $(C=O)_aO_bC_1-C_{10}$  alkyl,
- 3)  $(C=O)_aO_b$ aryl,
- 4)  $CO_2H$ ,
- 5) halo,
- 6)  $CN$ ,
- 7)  $OH$ ,
- 8)  $O_bC_1-C_6$  perfluoroalkyl,
- 9)  $O_a(C=O)_bNR^8R^9$ ,
- 10)  $S(O)_mR^a$ ,
- 11)  $S(O)_2NR^8R^9$ , and
- 12)  $-OPO(OH)_2$ ;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one,  
 15 two or three substituents selected from  $R^7$ ;

$R^6$  is independently selected from:

- 1)  $(C=O)_aO_bC_1-C_{10}$  alkyl,
- 2)  $(C=O)_aO_b$ aryl,
- 20 3)  $C_2-C_{10}$  alkenyl,
- 4)  $C_2-C_{10}$  alkynyl,
- 5)  $(C=O)_aO_b$  heterocyclyl,
- 6)  $CO_2H$ ,
- 7) halo,
- 25 8)  $CN$ ,
- 9)  $OH$ ,
- 10)  $O_bC_1-C_6$  perfluoroalkyl,
- 11)  $O_a(C=O)_bNR^8R^9$ ,
- 12)  $S(O)_mR^a$ ,
- 30 13)  $S(O)_2NR^8R^9$ ,
- 14) oxo,
- 15)  $CHO$ ,
- 16)  $(N=O)R^8R^9$ ,
- 17)  $(C=O)_aO_bC_3-C_8$  cycloalkyl, and
- 35 18)  $-OPO(OH)_2$ ;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

5 R<sup>7</sup> is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 3) oxo,
- 4) OH,
- 10 5) halo,
- 6) CN,
- 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 15 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,
- 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,
- 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>,
- 13) C(O)R<sup>a</sup>,
- 14) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>,
- 20 15) C(O)H,
- 16) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H, and
- 17) (C=O)<sub>r</sub>N(R<sup>b</sup>)<sub>2</sub>,
- 18) S(O)<sub>m</sub>R<sup>a</sup>,
- 19) S(O)<sub>2</sub>N(R<sup>b</sup>)<sub>2</sub>; and
- 25 20) -OPO(OH)<sub>2</sub>;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R<sup>b</sup>, OH, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, CO<sub>2</sub>H, CN, O(C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, oxo, NO<sub>2</sub> and N(R<sup>b</sup>)<sub>2</sub>;

30

R<sup>8</sup> and R<sup>9</sup> are independently selected from:

- 1) H,
- 2) (C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) (C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 35 4) (C=O)O<sub>b</sub>aryl,

- 5) (C=O)O<sup>b</sup>heterocyclyl,  
 6) C<sub>1</sub>-C<sub>10</sub> alkyl,  
 7) aryl,  
 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,  
 5 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,  
 10) heterocyclyl,  
 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 12) SO<sub>2</sub>R<sup>a</sup>, and  
 13) (C=O)NR<sup>b</sup><sub>2</sub>,
- 10 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>, or

R<sup>8</sup> and R<sup>9</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in  
 15 addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>10</sup> and R<sup>11</sup> are independently selected from: F and -CH<sub>2</sub>F;

20 R<sup>12</sup> and R<sup>13</sup> are independently selected from: H and -CH<sub>2</sub>F;

R<sup>ox</sup> is absent or is oxo;

25 R<sup>a</sup> is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>b</sup> is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>Re<sup>'</sup> or  
 30 S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, NH<sub>2</sub>, OH, OR<sup>a</sup>, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>Re<sup>'</sup>, S(O)<sub>2</sub>R<sup>a</sup> and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N(R<sup>b</sup>)<sub>2</sub>, wherein the  
 35 alkyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or


$R^c$  and  $R^{c'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said  
 5 monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^7$ ;

$R^d$  is selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, -(C<sub>2</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl and -  
 (C<sub>1</sub>-C<sub>6</sub>)alkyl-N( $R^b$ )<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three  
 10 substituents selected from  $R^7$ ;

$R^e$  and  $R^{e'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-  
 C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from  $R^7$ ; or

$R^e$  and  $R^{e'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said  
 15 monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^7$ .

2. The compound according to Claim 1 of Formula II:



or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

- a is 0 or 1;  
 5 b is 0 or 1;  
 m is 0, 1, or 2;  
 n is 0, 1, 2 or 3;  
 r is 0 or 1;  
 s is 0 or 1;  
 10 t is 0 or 1;

R<sup>1</sup> and R<sup>2</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

15 R<sup>3</sup> is selected from:

- 1) hydrogen;
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl;
- 3) C<sub>1</sub>-C<sub>10</sub> alkyl-O-R<sup>d</sup>,
- 4) C<sub>2</sub>-C<sub>10</sub> alkenyl-O-R<sup>d</sup>,
- 20 5) C<sub>2</sub>-C<sub>10</sub> alkynyl-O-R<sup>d</sup>,
- 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-O-R<sup>d</sup>,
- 7) C<sub>1</sub>-C<sub>10</sub> alkyl-(C=O)<sub>b</sub>-NR<sup>c</sup>R<sup>c'</sup>,
- 8) C<sub>2</sub>-C<sub>10</sub> alkenyl-(C=O)<sub>b</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 9) C<sub>2</sub>-C<sub>10</sub> alkynyl-(C=O)<sub>b</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 25 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl-(C=O)<sub>b</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 11) C<sub>1</sub>-C<sub>10</sub> alkyl-S(O)<sub>m</sub>-R<sup>d</sup>,
- 12) C<sub>2</sub>-C<sub>10</sub> alkenyl- S(O)<sub>m</sub>-R<sup>d</sup>,
- 13) C<sub>2</sub>-C<sub>10</sub> alkynyl- S(O)<sub>m</sub>-R<sup>d</sup>,
- 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl- S(O)<sub>m</sub>-R<sup>d</sup>,

30 said alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more substituents selected from R<sup>6</sup>;

R<sup>4</sup> is independently selected from:

- 35 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,



- 2)  $(\text{C}=\text{O})_a\text{O}_b\text{aryl}$ ,
- 3)  $\text{CO}_2\text{H}$ ,
- 4) halo,
- 5)  $\text{CN}$ ,
- 6)  $\text{OH}$ ,
- 7)  $\text{O}_b\text{C}_1\text{-C}_6$  perfluoroalkyl,
- 8)  $\text{O}_a(\text{C}=\text{O})_b\text{NR}^8\text{R}^9$ ,
- 9)  $\text{S}(\text{O})_m\text{R}^a$ ,
- 10)  $\text{S}(\text{O})_2\text{NR}^8\text{R}^9$ ,

10 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ;

$\text{R}^5$  is selected from:

- 1) hydrogen;
- 15 2)  $(\text{C}=\text{O})_a\text{O}_b\text{C}_1\text{-C}_{10}$  alkyl,
- 3)  $(\text{C}=\text{O})_a\text{O}_b\text{aryl}$ ,
- 4)  $\text{CO}_2\text{H}$ ,
- 5) halo,
- 6)  $\text{CN}$ ,
- 20 7)  $\text{OH}$ ,
- 8)  $\text{O}_b\text{C}_1\text{-C}_6$  perfluoroalkyl,
- 9)  $\text{O}_a(\text{C}=\text{O})_b\text{NR}^8\text{R}^9$ ,
- 10)  $\text{S}(\text{O})_m\text{R}^a$ ,
- 11)  $\text{S}(\text{O})_2\text{NR}^8\text{R}^9$ , and
- 25 12)  $-\text{OPO}(\text{OH})_2$ ;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ;

30  $\text{R}^6$  is independently selected from:

- 1)  $(\text{C}=\text{O})_a\text{O}_b\text{C}_1\text{-C}_{10}$  alkyl,
- 2)  $(\text{C}=\text{O})_a\text{O}_b\text{aryl}$ ,
- 3)  $\text{C}_2\text{-C}_{10}$  alkenyl,
- 4)  $\text{C}_2\text{-C}_{10}$  alkynyl,
- 35 5)  $(\text{C}=\text{O})_a\text{O}_b$  heterocyclyl,

- 6)  $\text{CO}_2\text{H}$ ,  
 7) halo,  
 8)  $\text{CN}$ ,  
 9)  $\text{OH}$ ,  
 5 10)  $\text{O}_b\text{C}_1\text{-C}_6$  perfluoroalkyl,  
 11)  $\text{O}_a(\text{C}=\text{O})_b\text{NR}^8\text{R}^9$ ,  
 12)  $\text{S}(\text{O})_m\text{R}^a$ ,  
 13)  $\text{S}(\text{O})_2\text{NR}^8\text{R}^9$ ,  
 14) oxo,  
 10 15)  $\text{CHO}$ ,  
 16)  $(\text{N}=\text{O})\text{R}^8\text{R}^9$ ,  
 17)  $(\text{C}=\text{O})_a\text{O}_b\text{C}_3\text{-C}_8$  cycloalkyl, and  
 18)  $-\text{OPO}(\text{OH})_2$ ;

15 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from  $\text{R}^7$ ;

$\text{R}^7$  is selected from:

- 1)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_1\text{-C}_{10})\text{alkyl}$ ,  
 20 2)  $\text{O}_r(\text{C}_1\text{-C}_3)\text{perfluoroalkyl}$ ,  
 3) oxo,  
 4)  $\text{OH}$ ,  
 5) halo,  
 6)  $\text{CN}$ ,  
 25 7)  $(\text{C}_2\text{-C}_{10})\text{alkenyl}$ ,  
 8)  $(\text{C}_2\text{-C}_{10})\text{alkynyl}$ ,  
 9)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_3\text{-C}_6)\text{cycloalkyl}$ ,  
 10)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-aryl}$ ,  
 11)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-heterocyclyl}$ ,  
 30 12)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-N}(\text{R}^b)_2$ ,  
 13)  $\text{C}(\text{O})\text{R}^a$ ,  
 14)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{R}^a$ ,  
 15)  $\text{C}(\text{O})\text{H}$ ,  
 16)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{H}$ ,  
 35 17)  $\text{C}(\text{O})\text{N}(\text{R}^b)_2$ ,

- 18)  $S(O)_m R^a$ ,
- 19)  $S(O)_2 N(R^b)_2$ ; and
- 20)  $-OPO(OH)_2$ ;

5 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $R^b$ , OH,  $(C_1-C_6)$ alkoxy, halogen,  $CO_2H$ , CN,  $O(C=O)C_1-C_6$  alkyl, oxo,  $NO_2$  and  $N(R^b)_2$ ;

$R^8$  and  $R^9$  are independently selected from:

- 1) H,
- 10 2)  $(C=O)O_b C_1-C_{10}$  alkyl,
- 3)  $(C=O)O_b C_3-C_8$  cycloalkyl,
- 4)  $(C=O)O_b$ aryl,
- 5)  $(C=O)O_b$ heterocyclyl,
- 6)  $C_1-C_{10}$  alkyl,
- 15 7) aryl,
- 8)  $C_2-C_{10}$  alkenyl,
- 9)  $C_2-C_{10}$  alkynyl,
- 10) heterocyclyl,
- 11)  $C_3-C_8$  cycloalkyl,
- 20 12)  $SO_2 R^a$ , and
- 13)  $(C=O)NR^b_2$ ,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from  $R^7$ , or

25  $R^8$  and  $R^9$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^7$ ;

30  $R^{10}$  is selected from: F and  $-CH_2F$ ;

$R^{13}$  is selected from: H and  $-CH_2F$ , provided that if t is 1,  $R^{13}$  is H;

35  $R^{ox}$  is absent or is oxo;

$R^a$  is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from  $R^7$ ;

- 5  $R^b$  is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup> or S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one, two or three substituents selected from  $R^7$ ;

- 10  $R^c$  and  $R^{c'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, NH<sub>2</sub>, OH, OR<sup>a</sup>, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup>, S(O)<sub>2</sub>R<sup>a</sup> and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N( $R^b$ )<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from  $R^7$ ; or


- 15  $R^c$  and  $R^{c'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^7$ ;

- 20  $R^d$  is selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, -(C<sub>2</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N( $R^b$ )<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from  $R^7$ ;

- 25  $R^e$  and  $R^{e'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from  $R^7$ ; or

- 30  $R^e$  and  $R^{e'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^7$ .

### 3. The compound according to Claim 2 of the Formula III:



or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

5

a is 0 or 1;

b is 0 or 1;

m is 0, 1, or 2;

n is 0, 1 or 2;

10 r is 0 or 1;

s is 0 or 1;

R<sup>1</sup> and R<sup>2</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

15

R<sup>4</sup> is independently selected from:

- 1) halo,
- 2) OH,
- 3) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

20

R<sup>5</sup> is selected from:

- 1) hydrogen,
- 2) halo,
- 3) OH,

4)  $O_bC_1-C_6$  perfluoroalkyl,

$R^7$  is selected from:

- |    |     |  |
|----|-----|--|
|    | 1)  | $(C=O)_rO_s(C_1-C_{10})$ alkyl,              |
| 5  | 2)  | $O_r(C_1-C_3)$ perfluoroalkyl,               |
|    | 3)  | oxo,   |
|    | 4)  | OH,  |
|    | 5)  | halo,  |
|    | 6)  | CN,  |
| 10 | 7)  | $(C_2-C_{10})$ alkenyl,                      |
|    | 8)  | $(C_2-C_{10})$ alkynyl,                      |
|    | 9)  | $(C=O)_rO_s(C_3-C_6)$ cycloalkyl,            |
|    | 10) | $(C=O)_rO_s(C_0-C_6)$ alkylene-aryl,         |
|    | 11) | $(C=O)_rO_s(C_0-C_6)$ alkylene-heterocyclyl, |
| 15 | 12) | $(C=O)_rO_s(C_0-C_6)$ alkylene- $N(R^b)_2$ , |
|    | 13) | $C(O)R^a$ ,                                  |
|    | 14) | $(C_0-C_6)$ alkylene- $CO_2R^a$ ,            |
|    | 15) | $C(O)H$ ,                                    |
|    | 16) | $(C_0-C_6)$ alkylene- $CO_2H$ , and          |
| 20 | 17) | $C(O)N(R^b)_2$ ,                             |
|    | 18) | $S(O)_mR^a$ , and                            |
|    | 19) | $S(O)_2N(R^b)_2$ ;                           |

25 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $R^b$ , OH,  $(C_1-C_6)$ alkoxy, halogen,  $CO_2H$ , CN,  $O(C=O)C_1-C_6$  alkyl, oxo,  $NO_2$  and  $N(R^b)_2$ ;

$R^8$  and  $R^9$  are independently selected from:

- |    |    |                               |
|----|----|-------------------------------|
|    | 1) | H,                            |
| 30 | 2) | $(C=O)O_bC_1-C_{10}$ alkyl,   |
|    | 3) | $(C=O)O_bC_3-C_8$ cycloalkyl, |
|    | 4) | $(C=O)O_b$ aryl,              |
|    | 5) | $(C=O)O_b$ heterocyclyl,      |
|    | 6) | $C_1-C_{10}$ alkyl,           |
| 35 | 7) | aryl,                         |

- 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 10) heterocyclyl,
- 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 5 12) SO<sub>2</sub>R<sup>a</sup>, and
- 13) (C=O)NR<sup>b</sup><sub>2</sub>,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>, or

- 10 R<sup>8</sup> and R<sup>9</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

- 15 R<sup>a</sup> is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

- 20 R<sup>b</sup> is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup> or S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;


- 25 R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, NH<sub>2</sub>, OH, OR<sup>a</sup>, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup>, S(O)<sub>2</sub>R<sup>a</sup> and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N(R<sup>b</sup>)<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or

- 30 R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

- 35 R<sup>e</sup> and R<sup>e'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or

$R^e$  and  $R^e'$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents  
 5 selected from  $R^7$ .

4. The compound according to Claim 3 of the Formula IV:



10 or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

a is 0 or 1;  
 15 b is 0 or 1;  
 m is 0, 1, or 2;  
 r is 0 or 1;  
 s is 0 or 1;

20  $R^1$  and  $R^2$  are independently selected from: H and  $(C_1-C_6)$ alkyl, optionally substituted with one, two or three substituents selected from  $R^7$ ;

$R^4$  is independently selected from:

1) halo,  
 25 2) OH,



3)  $O_bC_1-C_6$  perfluoroalkyl,

$R^7$  is selected from:

- 1)  $(C=O)_rO_s(C_1-C_{10})$ alkyl,
- 5 2)  $O_r(C_1-C_3)$ perfluoroalkyl,
- 3) oxo,
- 4) OH,
- 5) halo,
- 6) CN,
- 10 7)  $(C_2-C_{10})$ alkenyl,
- 8)  $(C_2-C_{10})$ alkynyl,
- 9)  $(C=O)_rO_s(C_3-C_6)$ cycloalkyl,
- 10)  $(C=O)_rO_s(C_0-C_6)$ alkylene-aryl,
- 11)  $(C=O)_rO_s(C_0-C_6)$ alkylene-heterocyclyl,
- 15 12)  $(C=O)_rO_s(C_0-C_6)$ alkylene- $N(R^b)_2$ ,
- 13)  $C(O)R^a$ ,
- 14)  $(C_0-C_6)$ alkylene- $CO_2R^a$ ,
- 15)  $C(O)H$ ,
- 16)  $(C_0-C_6)$ alkylene- $CO_2H$ , and
- 20 17)  $C(O)N(R^b)_2$ ,
- 18)  $S(O)_mR^a$ , and
- 19)  $S(O)_2N(R^b)_2$ ;

25 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $R^b$ , OH,  $(C_1-C_6)$ alkoxy, halogen,  $CO_2H$ , CN,  $O(C=O)C_1-C_6$  alkyl, oxo,  $NO_2$  and  $N(R^b)_2$ ;

$R^8$  and  $R^9$  are independently selected from:

- 1) H,
- 30 2)  $(C=O)O_bC_1-C_{10}$  alkyl,
- 3)  $(C=O)O_bC_3-C_8$  cycloalkyl,
- 4)  $(C=O)O_b$ aryl,
- 5)  $(C=O)O_b$ heterocyclyl,
- 6)  $C_1-C_{10}$  alkyl,
- 35 7) aryl,

- 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 10) heterocyclyl,
- 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 12) SO<sub>2</sub>R<sup>a</sup>, and
- 13) (C=O)NR<sup>b</sup><sub>2</sub>,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>, or

- 10 R<sup>8</sup> and R<sup>9</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

- 15 R<sup>a</sup> is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

- 20 R<sup>b</sup> is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup> or S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one, two or three substituents selected from R<sup>7</sup>;


- 25 R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, NH<sub>2</sub>, OH, OR<sup>a</sup>, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-OH, -(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR<sup>e</sup>R<sup>e'</sup>, S(O)<sub>2</sub>R<sup>a</sup> and -(C<sub>1</sub>-C<sub>6</sub>)alkyl-N(R<sup>b</sup>)<sub>2</sub>, wherein the alkyl is optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or

- 30 R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>7</sup>;

- 35 R<sup>e</sup> and R<sup>e'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>7</sup>; or

$R^e$  and  $R^{e'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents  
 5 selected from  $R^7$ .

5 The compound according to Claim 4 of the Formula V:



or a pharmaceutically acceptable salt or stereoisomer thereof,


10

wherein:

$R^1$  and  $R^2$  are independently selected from: H and  $(C_1-C_6)$ alkyl.

15

6. The compound according to Claim 2 of the formula VI:



or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

5

$R^1$  and  $R^2$  are independently selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

7. A compound selected from:

10 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

15


(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

20 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 5 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(2*R*,4*R*)-2-(fluoromethyl)-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 10 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(2*S*,4*S*)-2-(fluoromethyl)-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methyl-1-oxidopiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 15 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoropiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-isopropylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 20 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*S*)-3-fluoropiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 25 or a pharmaceutically acceptable salt thereof.


8. A compound which is:



(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

5 or a pharmaceutically acceptable salt thereof.


9. A compound which is:



10 (2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*S*,4*R*)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

or a pharmaceutically acceptable salt thereof.


15 10. A compound which is:



(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(2*R*,4*R*)-2-(fluoromethyl)-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide

5 or a pharmaceutically acceptable salt thereof.


11. A compound which is:










(2*S*)-4-(2,5-Difluorophenyl)-*N*-[(3*R*,4*S*)-3-fluoro-1-methylpiperidin-4-yl]-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide








or a pharmaceutically acceptable salt thereof.








12. The compound according to Claim 1 selected from:



















R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H

















R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
Me	Me	Me	F	H
Me		Me	F	H
Me		Me	F	H


R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H








R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H








$R_1$	$R_2$	$R_3$	$R_4$	$R_5$
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H








R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me	CH <sub>2</sub> OH	Me	Cl	H
Me	CH <sub>2</sub> OH	Me	Br	H
Me	CH <sub>2</sub> OH	Me	CN	H
Me	CH <sub>2</sub> OH	Me	Me	H
Me	CH <sub>2</sub> OH	Me	CF <sub>3</sub>	H
Me	CH <sub>2</sub> OH	Me	NO <sub>2</sub>	H
Me	CH <sub>2</sub> OH	Me	F	OH
Me	CH <sub>2</sub> OH	Me	F	NH <sub>2</sub>
Me	CH <sub>2</sub> OH	Me	F	F
Me	CH <sub>2</sub> OH	Me	F	SH



















R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H

















R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
	CH <sub>2</sub> OH	Me	F	H
Me	Me	Me	F	H
Me		Me	F	H
Me		Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H
Me	CH <sub>2</sub> OH		F	H

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
Me	CH <sub>2</sub> OH	Me	Cl	H
Me	CH <sub>2</sub> OH	Me	Br	H
Me	CH <sub>2</sub> OH	Me	CN	H
Me	CH <sub>2</sub> OH	Me	Me	H
Me	CH <sub>2</sub> OH	Me	CF <sub>3</sub>	H
Me	CH <sub>2</sub> OH	Me	NO <sub>2</sub>	H
Me	CH <sub>2</sub> OH	Me	F	OH
Me	CH <sub>2</sub> OH	Me	F	NH <sub>2</sub>
Me	CH <sub>2</sub> OH	Me	F	F
Me	CH <sub>2</sub> OH	Me	F	SH

or a pharmaceutically acceptable salt or stereoisomer thereof.

13. A pharmaceutical composition that is comprised of a compound in accordance with Claim 1 and a pharmaceutically acceptable carrier.

5

14. A method of treating or preventing cancer in a mammal in need of such treatment that is comprised of administering to said mammal a therapeutically effective amount of a compound of Claim 1.

10

15. A method of treating cancer or preventing cancer in accordance with Claim 14 wherein the cancer is selected from cancers of the brain, genitourinary tract, lymphatic system, stomach, larynx and lung.

15

16. A method of treating or preventing cancer in accordance with Claim 14 wherein the cancer is selected from histiocytic lymphoma, lung adenocarcinoma, small cell lung cancers, pancreatic cancer, glioblastomas and breast carcinoma.

17. A process for making a pharmaceutical composition which comprises combining a compound of Claim 1 with a pharmaceutically acceptable carrier.

20

18. The composition of Claim 13 further comprising a second compound selected from:

25

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor,
- 10) an angiogenesis inhibitor, and
- 11) a PPAR- $\gamma$  agonist,
- 12) a PPAR- $\delta$  agonists;

30

35

- 13) an inhibitor of cell proliferation and survival signaling,



- 14) an agent that interferes with a cell cycle checkpoint, and
- 15) an apoptosis inducing agent.

19. The composition of Claim 18, wherein the second compound is an  
5 angiogenesis inhibitor selected from the group consisting of a tyrosine kinase inhibitor, an  
inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an  
inhibitor of platelet derived growth factor, an MMP inhibitor, an integrin blocker, interferon- $\alpha$ ,  
interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole,  
combretastatin A-4, squalamine, 6-O-(chloroacetyl-carbonyl)-fumagillol, thalidomide,  
10 angiostatin, troponin-1, and an antibody to VEGF.

20. The composition according to Claim 13 further comprising a proteasome  
inhibitor.

15 21. The composition according to Claim 13 further comprising a aurora kinase  
inhibitor.

22. The composition according to Claim 13 further comprising a Raf kinase  
inhibitor.

20 23. The composition according to Claim 13 further comprising a  
serine/threonine kinase inhibitor.

24. The composition according to Claim 13 further comprising an inhibitor of  
25 another mitotic kinesin which is not KSP.

25. The composition of Claim 19, wherein the second compound is an  
estrogen receptor modulator selected from tamoxifen and raloxifene.

30 26. A method of treating cancer which comprises administering a  
therapeutically effective amount of a compound of Claim 1 in combination with radiation  
therapy.

27. A method of treating or preventing cancer that comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a compound selected from:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor,
- 10) an angiogenesis inhibitor,
- 11) PPAR- $\gamma$  agonists,
- 12) PPAR- $\delta$  agonists,
- 13) an inhibitor of inherent multidrug resistance,
- 14) an anti-emetic agent,
- 15) an agent useful in the treatment of anemia,
- 16) an agent useful in the treatment of neutropenia,
- 17) an immunologic-enhancing drug,
- 18) an inhibitor of cell proliferation and survival signaling,
- 19) an agent that interferes with a cell cycle checkpoint, and
- 20) an apoptosis inducing agent.

28. A method of treating cancer that comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with radiation therapy and a compound selected from:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,

- 5                    9)      a reverse transcriptase inhibitor,  
                      10)     an angiogenesis inhibitor,  
                      11)     PPAR- $\gamma$  agonists,  
                      12)     PPAR- $\delta$  agonists,  
                      13)     an inhibitor of inherent multidrug resistance,  
                      14)     an anti-emetic agent,  
                      15)     an agent useful in the treatment of anemia,  
                      16)     an agent useful in the treatment of neutropenia,  
                      17)     an immunologic-enhancing drug,  
 10                  18)     an inhibitor of cell proliferation and survival signaling,  
                      19)     an agent that interferes with a cell cycle checkpoint, and  
                      20)     an apoptosis inducing agent.

15                  29.      A method of treating or preventing cancer which comprises administering  
 a therapeutically effective amount of a compound of Claim 1 and paclitaxel or trastuzumab.

                     30.      A method of treating or preventing cancer which comprises administering  
 a therapeutically effective amount of a compound of Claim 1 and a GPIIb/IIIa antagonist.

20                  31.      The method of Claim 30 wherein the GPIIb/IIIa antagonist is tirofiban.

                     32.      A method of treating or preventing cancer which comprises administering  
 a therapeutically effective amount of a compound of Claim 1 in combination with a COX-2  
 inhibitor.

25                  33.      A method of treating or preventing cancer which comprises administering  
 a therapeutically effective amount of a compound of Claim 1 in combination with a proteasome  
 inhibitor.

30                  34.      A method of treating or preventing cancer which comprises administering  
 a therapeutically effective amount of a compound of Claim 1 in combination with an aurora  
 kinase inhibitor.

35. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a Raf kinase inhibitor.

5 36. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a serine/threonine kinase inhibitor.

10 37. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with an inhibitor of a mitotic kinesin that is not KSP.

15 38. A method of modulating mitotic spindle formation which comprises administering a therapeutically effective amount of a compound of Claim 1.

39. A method of inhibiting the mitotic kinesin KSP which comprises administering a therapeutically effective amount of a compound of Claim 1.

**TITLE OF THE INVENTION**

**MITOTIC KINESIN INHIBITORS**

**ABSTRACT OF THE DISCLOSURE**

5           The present invention relates to dihydropyrrole compounds that are useful for treating cellular proliferative diseases, for treating disorders associated with KSP kinesin activity, and for inhibiting KSP kinesin. The invention is also related to compositions which comprise these compounds, and methods of using them to treat cancer in mammals.

10

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/026012

International filing date: 11 August 2004 (11.08.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/495,637  
Filing date: 15 August 2003 (15.08.2003)

Date of receipt at the International Bureau: 27 September 2004 (27.09.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse